



**Report on Epidemiological Mapping of Schistosomiasis and
Soil Transmitted Helminthiasis in 19 States and the FCT,
Nigeria.**



May, 2015



Report on Epidemiological Mapping of Schistosomiasis and Soil Transmitted Helminthiasis in 19 States and the FCT, Nigeria.



TRAINING SITE - 'STATE A'



Welcome to the project site for the Epidemiological Mapping of Schistosomiasis and Soil Transmitted Helminthiasis (STH) in Nigeria. This website is restricted to authorized users only.

Survey Datasets

File is downloaded as .xls (Microsoft Excel)

These data are password protected and not accessible to the general public or others without the proper account information.

- [School survey \(53\)](#)
- [Students survey \(1672\)](#)
- [Stool survey \(1422\)](#)
- [Urine survey \(2350\)](#)
- [Error survey \(33\)](#)
- [Download linked dataset](#)

Coding guides:

- [School](#)
- [Students](#)

Progress Reports

| Survey Name | Last Submission | # Total Records |
|-------------|---------------------|-----------------|
| School | 2014-02-23 15:12:07 | 53 |
| Students | 2014-02-23 15:10:06 | 1672 |
| Stool | 2014-02-23 15:10:14 | 1422 |
| Urine | 2014-02-23 15:10:26 | 2350 |
| Error | 2014-02-19 08:49:20 | 33 |

Student Survey

Viewing last 10 entries only

| Submitted | Community/school | Individuals |
|------------|------------------|-------------|
| 2014-02-14 | 1887 | 50 |
| 2014-02-13 | 1881 | 48 |
| 2014-02-13 | 1895 | 48 |
| 2014-02-13 | 1886 | 50 |
| 2014-02-13 | 1891 | 48 |

TABLE OF CONTENTS

| | |
|---|-----|
| LIST OF FIGURES | v |
| LIST OF PLATES | vii |
| FOREWORD..... | x |
| EXECUTIVE SUMMARY..... | xii |
| 1.0 BACKGROUND | 1 |
| 1.1 Introduction..... | 1 |
| 1.2 Objectives of the Mapping Project..... | 2 |
| 1.3 Justification for the Survey..... | 2 |
| 2.0. MAPPING METHODOLOGY..... | 3 |
| 2.1 Study Area | 3 |
| 2.2. Study Design | 4 |
| 2.2.1 Selection of Participating Schools and Children | 4 |
| 2.3 School and Community Mobilization and Sensitization..... | 5 |
| 2.4 Training of Personnel..... | 5 |
| 2.5 Field Data Collection Process and Uploading into Cloud Server | 8 |
| 2.6 Sample Collection and Examination | 9 |
| 2.7 Data Analyses | 10 |
| 2.8 Ethical Clearance | 11 |
| 3.0 RESULTS | 12 |
| 3.1 Demographic data | 12 |
| 3.2 Parasitological Findings | 12 |
| 3.1.1 Schistosomiasis Result | 16 |
| 3.1.2 Soil Transmitted Helminths (STHs) Results..... | 19 |
| 3.1.3 Prevalence of schistosomiasis and STH by Sex | 22 |
| 3.1.4 Prevalence of schistosomiasis and STH by Age group | 24 |
| 3.1.5 Intensity of Infection..... | 26 |
| 3.2 Implication of Findings on Intervention | 27 |
| 3.3 Defecation practices and water contact activities | 31 |

| | |
|---|----|
| 4.0 DISCUSSIONS | 33 |
| 4.1 Discussion of Results | 33 |
| 4.2 Implications of Findings..... | 35 |
| 4.3 Some Key Achievements of the Project | 36 |
| 5.0. CHALLENGES/CONSTRAINTS | 37 |
| 5.1 Logistics | 37 |
| 5.2 Mobilisation/Misconception | 37 |
| 5.3 Communication/Technology Challenges..... | 37 |
| 5.4 Access challenges to schools in some LGAs | 37 |
| 5.5 Security Challenges..... | 38 |
| 6.0. RECOMMENDATIONS | 39 |
| REFERENCES | 41 |
| APPENDIX | 43 |
| Appendix 1: Coordinated Mapping of Schistosomiasis & STH in Nigeria: Project Organogram | 43 |
| Appendix 2: Survey Teams | 44 |
| Appendix 3: Data collection tools..... | 49 |
| Appendix 4: Flow Diagram of the Android Based Electronic Data Management | 58 |
| Appendix 5: Ethical Clearance | 59 |
| Appendix 6: Treatment guideline for schistosomiasis/STH..... | 60 |
| Appendix 7: LGAs Requiring Interventions..... | 61 |
| Appendix 8: Schistosomiasis and STHs Prevalence by Practice of Defecation..... | 72 |
| Appendix 9: Prevalence of Schistosomiasis and STHs by water contact activities | 75 |

LIST OF TABLES

| | |
|---|----|
| Table 1: Categories of Personnel Trained for the Survey..... | 6 |
| Table 2: Demographic Characteristics of the Study Population..... | 12 |
| Table 3: Prevalence of Schistosomiasis, Soil Transmitted Helminths and Co-Infection..... | 13 |
| Table 4: Range of Schistosomiasis and STHs Prevalence by State | 15 |
| Table 5: Parasite intensity in eggs per gram (epg) of faeces or per 10 ml of urine | 27 |
| Table 6: LGAs requiring Intervention for Schistosomiasis..... | 28 |
| Table 7: LGAs requiring Intervention for Soil Transmitted Helminths (STHs)..... | 30 |
| Table 8: Prevalence of schistosomiasis and STHs in relation to places of defecation. | 31 |
| Table 9: Prevalence of Schistosomiasis in relation to water contact activities..... | 32 |

LIST OF FIGURES

| | |
|---|-----------|
| Figure 1: Map of Nigeria showing study area..... | 3 |
| <i>Figure 2: Map of Nigeria showing the study area and funding partners</i> | <i>4</i> |
| <i>Figure 3: Schistosomiasis and STHs endemic areas.....</i> | <i>14</i> |
| <i>Figure 4: Prevalence of Schistosomiasis by LGA</i> | <i>15</i> |
| <i>Figure 5: Prevalence of STHs by LGA</i> | <i>16</i> |
| <i>Figure 6: Proportion of S. haematobium and S. mansoni in the study area.....</i> | <i>16</i> |
| Figure 7: Prevalence of S. haematobium and S.mansoni by State..... | 17 |
| Figure 8: Schistosomiasis Point Prevalence maps | 18 |
| <i>Figure 9: Proportion of the three species of STHs in the Study Area</i> | <i>19</i> |
| <i>Figure 10: Prevalence of A. lumbricoides, Hookworm and T. trichiura</i> | <i>20</i> |
| <i>Figure 11: STHs Point Prevalence maps.....</i> | <i>21</i> |
| <i>Figure 12: Proportion of pupils infected with Schistosomiasis by sex</i> | <i>22</i> |
| <i>Figure 13: Prevalence of Schistosomiasis by sex</i> | <i>22</i> |
| <i>Figure 14: Proportion of Pupils infected with STHs by sex.....</i> | <i>23</i> |
| <i>Figure 15: Prevalence of STHs by sex.....</i> | <i>23</i> |
| <i>Figure 16: Proportion of schistosomiasis infected pupils by age group</i> | <i>24</i> |
| <i>Figure 17: Prevalence of schistosomiasis by age group</i> | <i>25</i> |
| <i>Figure 18: Proportion of STHs infected pupils by age group</i> | <i>25</i> |

Figure 19: Prevalence of STHs infected pupils by age group 26

Figure 20: Intervention planning map for Schistosomiasis in 19 States and FCT. 29

Figure 21: Intervention planning map for STHs in 19 States and the FCT 31

LIST OF PLATES

Plates 1a & 1b: A cross section of school pupils being selected for sample collection 5

Plate 2a: *Advocacy visit to Education Secretary.* 5
Plate 2b: *Advocacy visit to the Hon. Commissioner for Health*..... 5

Plate 3: *Participants and Facilitators for the Quantum GIS training at Abuja* 7

Plate 4: *Laboratory training session.* 7

Plate 5: *State survey team*..... 8

Plate 6: *A Recorder inputting student’s information to the android phone* 9

Plate 7: *Water contact behavioural activities* 9

Plate 8: *Technicians Preparing Kato-Katz in the laboratory.*..... 10

Plate 9: *Technicians conducting urinalysis in the laboratory.* 10

ACRONYMS

| | |
|----------|---|
| CDC | Centers for Disease Control and Prevention |
| CIFF | Children's Investment Fund Foundation |
| DFID | Department for International Development |
| FCT | Federal Capital Territory |
| FMOH | Federal Ministry of Health |
| GTMP | Global Trachoma Mapping Project |
| HANDS | Health and Development Support |
| HKI | Helen Keller International |
| IBM | International Business Machine |
| LGA | Local Government Area |
| LSHTM | London School of Hygiene and Tropical Medicine |
| MAM | Mass Administration of Medicines |
| MITOSATH | Mission to Save the Helpless |
| NBS | National Bureau of Statistics |
| NGDOs | Non-Governmental Development Organizations |
| NHMIS | National Health Management Information System |
| NHREC | National Health Research Ethical Committee |
| NPC | National Population Commission |
| NTDs | Neglected Tropical Diseases |
| PAG | Project Advisory Group |
| PCT | Preventive Chemotherapy |
| PHASE | Preventive chemotherapy, Access to clean water, Sanitation improvement and environmental manipulation |
| QGIS | Quantum Geographic Information System |
| SCI | Schistosomiasis Control Initiative |
| SDGs | Sustainable Development Goals |
| SMOH | State Ministry of Health |
| SPSS | Statistical Package for Social Science |
| STHs | Soil Transmitted Helminths |

| | |
|-------|---------------------------------------|
| SUBEB | State Universal Basic Education Board |
| TCC | The Carter Center |
| TFGH | Task Force for Global Health |
| UK | United Kingdom |
| WASH | Water, Sanitation and Hygiene |
| WHO | World Health Organization |

FOREWORD

Schistosomiasis and Soil Transmitted Helminths (STHs) are among the group of Neglected Tropical Diseases (NTDs) that cause chronic infections and ill health in endemic areas. Nigeria is one of the countries that have the highest burden of these NTDs. In the last two years, Nigeria has scaled-up epidemiological mapping for NTDs with the support of partners and stakeholders. There are further plans to scale up Preventive Chemotherapy, Health Education, Access to clean water, Sanitation improvement and Environmental manipulation (PHASE) activities to reduce disease burden.

This report provides epidemiological information on 19 States and the Federal Capital Territory (FCT) that were mapped for schistosomiasis and STHs from November 2013 to May 2015. The Children's Investment Fund Foundation (CIFF) provided funding to map 14 States and the remaining States were supported by Department for International Development (DFID) funded Global Trachoma Mapping Project (GTMP), RTI/ENVISION project and Sightsavers. Findings from the survey showed that all 19 States and the FCT are endemic for schistosomiasis or STHs and in some cases both. This provides vital evidence for improved strategic planning for schistosomiasis and STHs integrated disease control and elimination in Nigeria.

One of the key innovations of this project was transition from use of paper-based questionnaires to electronic data collection tools, the Android-based smart phones and cloud server platform. The electronic-based mapping project provided opportunity for enhanced data collection, timely transmission and reporting as well as data security. The completion of the Schistosomiasis and STHs epidemiological mapping in 19 States and the FCT is an important milestone in the fight against NTDs and has given Nigeria a pride of place in the global NTDs elimination landscape.

I wish to convey the nation's appreciation to the funding organizations and other stakeholders that have worked tirelessly to achieve this feat. I would also like to remind everyone that this is a giant stride in our collective and collaborative effort towards meeting the national and global NTDs elimination goals. The findings from this project provide evidence-based data for decision making in order to appropriately target interventions. It is expected that our children at risk will benefit immensely from this for a brighter future.

I wish to reaffirm the commitment of the Federal Government of Nigeria to the elimination of NTDs in Nigeria and in particular schistosomiasis and STHs.



Engr. Fidelis. N. Nwankwo
Honourable Minister of State for Health

ACKNOWLEDGEMENTS

The Federal Ministry of Health sincerely acknowledges the immense support and contributions of our NGDO partners especially Sightsavers, HANDS, HKI, TCC and MITOSATH towards the control and elimination of Schistosomiasis and STHs in Nigeria.

Special appreciation goes to the Children's Investment Fund Foundation (CIFF) and partners; Sightsavers, RTI/ENVISION and DFID for their tremendous financial and technical support and humanitarian gesture to conduct the Schistosomiasis and STHs epidemiological mapping in 19 States and the FCT.

We also thank the Project Advisory Group (PAG) and Schistosomiasis Control Initiative (SCI), UK for their advice and useful insight towards the development of the mapping protocol and survey tools.

The Honourable Commissioners for Health in the States and the FCT Secretary for Health are appreciated for their collaboration and support during the mapping activities. We also recognize the effort and contributions of the consultants, technical officers from the Federal and State Ministries of Health and other support personnel at all levels of operation. The management and staff of the 19 States and FCT, State Universal Basic Education Board (SUBEB), the LGAs, participating schools and members of the surveyed communities are most appreciated for their support and cooperation throughout the duration of the exercise.

Our immense thanks go to the school children for their cooperation and enthusiastic participation in the mapping activities.



Dr Bridget Okeoguale
Director, Public Health

EXECUTIVE SUMMARY

Epidemiological mapping is a prerequisite for Neglected Tropical Diseases (NTDs) Preventive chemotherapy, Access to clean water, Sanitation improvement and Environmental manipulation (PHASE) interventions.

The Children's Investment Fund Foundation (CIFF) as part of its investment in Nigeria provided funding to map schistosomiasis and Soil Transmitted Helminths (STHs) in 347 LGAs in 14 States. This support provided leverage for additional funding to map 86 LGAs in five other States and the Federal Capital Territory (FCT). The coordinated mapping of schistosomiasis and STHs was conducted in 19 States and the FCT of Nigeria from November 2013 to May 2015.

A sample of 50 - 55 children from five randomly selected schools in each of the 433 LGAs in 19 States and four Area Councils of the FCT were examined for schistosomiasis and STHs with generous funding support from CIFF, Sightsavers, Department for International Development (DFID) funded Global Trachoma Mapping Project (GTMP) and the RTI/ENVISION project.

Epidemiological data on both diseases were mapped using a novel technique; the LINKS system developed by the Task Force for Global Health (TFGH). This system uses a collection of open source tools for data collection on Android devices and cloud based data reporting and management. The application of these devices supported the transition from paper-based questionnaires to electronic data collection tools. Also, Water, Sanitation and Hygiene (WASH) information for schools and school children were collected. The Kato-Katz technique, dipsticks (Haemastix), syringe filtration and sedimentation techniques were used to examine stool and urine samples collected from the school children.

The result of this survey revealed that all the States and the FCT were endemic for one or both diseases with an overall prevalence of 9.5% for schistosomiasis and 27% for STHs suggesting low risks of both diseases in the project area. However, the data captured by LGA; the intervention unit showed that prevalence of infections varied from low to high risk. Of the 433 LGAs surveyed the number of LGAs requiring interventions for schistosomiasis and STHs were 359 and 237 respectively. The prevalence of infection was significantly higher in males than in females for both diseases. STHs were more prevalent among the younger age group (5-10years) while schistosomiasis was more prevalent among the older age group (11-16 years). Two percent of the pupils surveyed were co-infected with schistosomes and STHs.

Schistosomiasis and STHs were seen among pupils who claimed to defecate in the school toilets, around the school compound and outside school environment. The mapping exercise provided insight into disease distribution and intensity in the 19 States and FCT surveyed.

At the end of the exercise, the capacities of the national and state government personnel have been strengthened in implementing and evaluating planned NTD control activities effectively. Working with the various partners has provided a platform for cross learning, skills sharing and ultimately improved programme coordination.

There were several engagements with the Honorable Commissioners for Health and other stakeholders to advocate for support to the project, present the findings and need for rapid public health action. At the community level awareness on the prevalence of schistosomiasis and STHs were raised to secure their participation.

The mapping of schistosomiasis and STHs in 19 States and the Federal Capital Territory of Nigeria was successfully completed and has achieved all the set objectives. Data generated from this crucial survey provided vital evidence for appropriate and sustainable intervention by Government in collaboration with our highly esteemed NTDs partners. It is the right of every child in Nigeria to enjoy good health and the time to intervene is now.

It is recommended that Governments (Federal, State and LGAs) and stakeholders scale up uninterrupted provision and administration of appropriate medicines which should be implemented alongside other interventions in the PHASE strategy. Plans should be put in place for impact assessment after the third year of consistent Mass Administration of Medicines (MAM).

1.0 BACKGROUND

1.1 Introduction

Schistosomiasis and Soil Transmitted Helminths (STHs) are Neglected Tropical Diseases (NTD) that cause ill-health and chronic infections in endemic countries, including Nigeria. The country is estimated to have the highest number of people infected with NTDs in Africa; a group of parasitic and bacterial infections that affect the world's poorest populations (WHO 2015; Hotez and Kamath 2009). Nigeria has the highest burden of endemicity of intestinal helminth infections and cases of schistosomiasis. The latter includes intestinal schistosomiasis caused by *Schistosoma mansoni* and urogenital schistosomiasis caused by *Schistosoma haematobium* (WHO 2015; Hotez and Kamath 2009). These parasites are known to have detrimental impact on child health as they deplete nutrients in children and adversely affect physical and cognitive development, causing symptoms such as abdominal pain, anaemia, bladder and liver diseases and other health problems which impair growth, reduce school attendance with poor learning outcomes (Lobato *et al.*, 2012).

The exact national distribution and disease burden of schistosomiasis and STHs was largely unknown before the current scale up of epidemiological mapping activities in Nigeria. The national data available were from limited number of epidemiological surveys conducted between 2008 and 2012, among school age children in 207 of the 774 LGAs. Although mapping had not been conducted in all 36 States and the FCT, schistosomiasis and STHs were suspected to be co-endemic in majority of the States.

As part of the ongoing effort to scale up epidemiological mapping activities in Nigeria, the Federal Ministry of Health (FMOH), in collaboration with various partners and stakeholders launched several disease mapping projects nationwide (Appendix 1). Within this effort CIFF provided Eight hundred and eighty nine thousand, five hundred and sixty three pounds sterling (£889,563.00) to map schistosomiasis and STHs in 347 LGAs in 14 States. This support enabled FMOH to standardise a training manual for schistosomiasis and STHs mapping that eased data collection and management. This also led to the development of national mapping protocol and data collection platform used by other donors to support the field data collection. The national protocol was used as a standard and reference document during training of the field personnel and throughout the mapping process. The technical support from Schistosomiasis Control Initiative (SCI) provided further guidance and insight into the design and development of these mapping tools. Other partners leveraged on the support provided by CIFF to map other States. The DFID-funded GTMP mapped Bauchi, Niger and Taraba States; RTI/ENVISION mapped Cross River and Ondo States while Sightsavers mapped Benue State. Technical officers from FMOH, in-country consultants and NGDO partners were trained on the use of

Geographic Information System (GIS). The skills acquired were used to capture, store, analyse and manage spatial data and also to develop disease and intervention maps.

1.2 Objectives of the Mapping Project

- i. To complete epidemiological mapping of schistosomiasis and STHs in the 347 LGAs in 14 States of Nigeria over an 18 month period
- ii. To provide the necessary evidence of disease burden to be used for advocacy to leverage government funding and inform the broader NTD community beyond Nigeria
- iii. To develop the capacity of government personnel on epidemiological mapping for schistosomiasis and STHs, and ensure commitment to implement a national deworming programme

1.3 Justification for the Survey

Schistosomiasis and STHs remain a serious public health problem in Nigeria. Assessment of disease prevalence and the geographic distribution are the first steps in planning effective intervention measures for schistosomiasis and STHs.

The completion of epidemiological mapping equally presented a unique opportunity to collect for the first time, a more comprehensive data on the national disease distribution. Specific national information on the prevalence, distribution and disease burden resulting from these NTDs provided a basis for prioritizing control programme strategies in line with the national NTDs Master Plan and the Global Elimination target. The findings of the survey are a useful advocacy tool for leveraging Government resources and accessing global donated NTD medicines. Comprehensive up-to-date maps showing the diseases distribution will facilitate efficient resource utilization and appropriate drugs distribution to target populations.

The benefits of control and elimination of these diseases include reduction of school absenteeism, malnutrition and anaemia leading to improvement of growth, physical and intellectual development and attainment of full potentials by children in Nigeria. It will also contribute to poverty reduction and improved socio-economic wellbeing of families. Giving the strong association between helminthic, other NTDs, economic and human development, it has become essential to give due consideration to eliminating helminths infection as a means to achieve the Sustainable Development Goals (SDGs) in Nigeria (Hotez and Herricks 2015).

2.0. MAPPING METHODOLOGY

2.1 Study Area

Nigeria is the most populous country in Africa with a projected population of over 170 million, National Population Commission (NPC 2006). Nigeria occupies the area along the West Coast of Africa between latitude 4^o and 14^o N and longitude 5^o and 14^o E covering about 923,768 square kilometres and bordered in the North by the Republics of Niger and Chad and in the East by Cameroon while Benin Republic borders it on the West and Atlantic Ocean to the south. Nigeria has a coast line of about 3,122 kilometres. The capital city Abuja is located within the Federal Capital Territory, which is about 713 square kilometres. There are 36 States and the FCT with 774 Local Government Areas (LGAs) in the Federation.

The study area comprised 19 States and the FCT as shown in Figure 1. Altogether the 19 States and the FCT have 456 LGAs and an estimated population of 106,243,198 projected 2006 population (NPC 2006). The survey was conducted in 433 out of the 456 LGAs, 21 were previously mapped and two were not mapped due to security challenges. The Children's Investment Fund Foundation and other partners provided funding support for mapping activities in different States (Figure 2).




Figure 1: Map of Nigeria showing study area




Figure 2: Map of Nigeria showing the study area and funding partners

2.2. Study Design

State-wide mapping for schistosomiasis and STHs infections was conducted in a coordinated manner using National protocol based on WHO (2010) framework. The survey was cross sectional and purposive at State and LGA levels aimed at completing epidemiological mapping in Nigeria. There was a randomised selection of schools from the sampling frame followed by a randomised systematic selection of children in the schools surveyed. The sampling frame was the list of all primary schools in each ward. A stratified random cluster sampling procedure was used in line with the FMOH protocol on integrated epidemiological mapping and baseline survey for schistosomiasis and STHs (FMOH 2013). Primary schools in northern Nigeria included non-formal schools (“Madrasat”/Islamic Schools).

The survey was based on standard diagnostic procedures for collection and examination of urine and faecal samples from school age children for the presence of schistosome and intestinal helminth eggs. Enrolled school age children were targeted from the surveyed communities. Males and females were selected on pro rata basis.

2.2.1 Selection of Participating Schools and Children

In all the LGAs of the States surveyed, five schools were randomly selected from different communities; however, schools in areas with large water bodies were prioritised. A sampling frame

was developed and used for selection of pupils in each selected school. A range of 50-55 pupils of both sexes from 5 – 16 years old from each school was sampled (Plates 1a & 1b).



Plates 1a & 1b: A cross section of school pupils being selected for sample collection

2.3 School and Community Mobilization and Sensitization

Advocacy visits were paid to the Honourable Commissioners for Health and this was preceded by letters from the FMOH. The State Universal Basic Education Board (SUBEB) personnel mobilised the schools. Letters were also sent from the States Ministry of Health through SUBEB to selected schools and communities. Some States also mobilised the communities through the mass media, including radio and television. Education secretaries, community health educators and districts heads in some communities were also sensitized (Plates 2a & 2b).



Plate 2a: Advocacy visit to Education Secretary.



Plate 2b: Advocacy visit to the Hon. Commissioner for Health

2.4 Training of Personnel

Training of personnel covered the use of the following:

- i. Electronic data collection device (LINKS system)
- ii. Backup forms (for data verification and validation)
- iii. Quantum Geographic Information System (QGIS)
- iv. Kato-Katz techniques (for stool analysis)
- v. Filtration and sedimentation technique (for urinalysis)

The training was conducted at National, Zonal and State levels. A total of 774 personnel were trained and participated in the mapping exercise (Table 1). A list of the survey team is in Appendix 2.

Table 1: Categories of Personnel Trained for the Survey

| S/N | Categories of personnel | Number trained |
|-----|-------------------------|----------------|
| 1 | Laboratory scientists | 100 |
| 2 | Laboratory technicians | 100 |
| 3 | Recorders | 100 |
| 4 | LGA Coordinators | 433 |
| 5 | In-country consultants | 8 |
| 6 | NGDO and FMOH personnel | 33 |
| | Total | 774 |

Trainings at the national level were on the use of LINKS mobile system and QGIS. The LINKS mobile system was developed by the Task Force for Global Health. The system uses a collection of open source tools for data collection on Android powered devices and cloud based data reporting and management. During the training session, the system was field tested and feedback given to the developer which facilitated necessary adjustment on the platform before commencement of project.

The QGIS training was facilitated by a team from the London School of Hygiene and Tropical Medicine (LSHTM), UK. This provided participants with an overview on the use of epidemiological mapping tools to assist in the implementation and evaluation of planned NTD control activities in Nigeria. Participants included FMOH staff, NGDOs, National Bureau of Statistics (NBS) and in-country consultants (Plate 3).

The Training of Trainers (ToT) at zonal level took place in four locations, two in the North (Jigawa and Bauchi States) and two in the South (Lagos and Ekiti States). The ToT in Jigawa State was followed by flag off of the survey by the Honourable Minister of Health. At the flag off event, participants included the State officials: the Honourable Commissioner for Health, Jigawa, trainers of trainers, NGDOs, in-country consultants and FMOH team.



Plate 3: Participants and Facilitators for the Quantum GIS training at Abuja

A training was cascaded for the field teams; comprising staff of State Ministries of Health and Education (including SUBEB) on the mapping methodology and community mobilization. Laboratory staff were trained on sample collection and examination while the recorders were trained on the use of electronic data capturing devices (Plates 4).



Plate 4: Laboratory training session.

Practical sessions and post training tests were also conducted. At the end of each State’s training, field teams were selected based on post-training performance. Micro planning meetings were held to discuss schools and community mobilization, survey approach and detailed implementation plans developed.

In each State, a team was constituted of recorders, scientists, technicians, State NTD programme officer, SUBEB representative, in-country consultant, FMOH supervisor and NGDO technical officer (Plate 5). This team was further divided into five sub-teams made up of a recorder, laboratory technician, scientist, a supervisor, a driver and a local guide.



Plate 5: State survey team

2.5 Field Data Collection Process and Uploading into Cloud Server

The electronic data collection tool has five forms; school and child information forms, urine and stool results, and error reporting forms (Appendix 3). These forms were downloaded into the android-based phones used by recorders for data collection in the field and transmission to the cloud server platform (Plate 6 and Appendix 4). Geographical coordinates of each sampled school and community were captured within the school premises in the process of collecting the school information. In order to document the knowledge, attitude and practices of the surveyed population as relates to disease transmission; Water, Sanitation and Hygiene (WASH) parameters were collected. Some of these are hand washing practices and availability of toilet facilities in the schools. Toilet facilities were ascertained and inspected.

The proximity of water bodies to schools and water contact behavioural activities (Plate 7) were documented in all the schools visited through interviews with school Heads and the pupils.



Plate 6: A Recorder inputting student's information to the android phone



Plate 7: Water contact behavioural activities

2.6 Sample Collection and Examination

Midstream urine and stool samples were collected from selected school children using sterile specimen bottles. Physical appearance of the urine samples collected ranged from clear, amber, pale, cloudy and bloody. These samples were tested for haematuria using Combi-9[®] reagent strip and were examined in the laboratory for schistosome eggs using urine filtration/sedimentation technique (Lengeler *et al.*, 1993). Stool samples collected from the field were examined for parasite eggs using the Kato-Katz technique (WHO,1991; 2010), as shown in Plates 8 and 9. Quality control was instituted

and laboratory results were verified by the consultants and FMOH supervisors to ensure consistency in samples preparation and examination.



Plate 8: Technicians Preparing Kato-Katz in the laboratory.



Plate 9: Technicians conducting urinalysis in the laboratory.

2.7 Data Analyses

Data cleaning by FMOH and NGOs data managers/technical officers was carried on completion of the mapping activities using specified guidelines. State specific linked datasets from the cloud server were downloaded and cleaned using excel add-in (Ablebit®). This was to validate any observed discrepancy between uploaded data and entries on back-up forms. Statistical analyses were carried out using IBM SPSS® version 20 and Epi Info 7.

Descriptive statistics such as percentages were calculated. Frequencies and prevalence of key indicators were generated and presented in tables, graphs, charts and maps. Chi-square test was used to investigate associations between categorical variables while Z-test for proportions was to compare proportions of diseases by gender and age groups. The statistical tests were carried out at 5% level of significance.

2.8 Ethical Clearance

Ethical clearance was obtained from the National Health Research Ethical Committee (NHREC) of the FMOH (Appendix 5). Ethical permissions obtained from the SMOH and SUBEB were conveyed to the schools. Head of schools acknowledged receipt and gave consent for the exercise to be carried out.

3.0 RESULTS

3.1 Demographic data

Schistosomiasis and STHs epidemiological survey was conducted in Nigeria between November, 2013 and May, 2015 in 19 States and the FCT in 433 unmapped LGAs covering 2,160 schools and communities. Samples were collected from 108,472 pupils comprising 57,670 (53.2%) males and 50,802(46.8%) females. The age range of the sampled pupils were between 5-10 years (57,599) and 11-16 years (50,873) Table 2.

Table 2: Demographic Characteristics of the Study Population

| State | No Surveyed | | | | | |
|-------------|--------------------------|----------------------|---------------|---------------|---------------|---------------|
| | LGAs (Surveyed/Total) | School/ Community | Female | Male | 5 - 10 Years | 11 - 16 Years |
| Akwa Ibom | 31 /31 | 155 | 3,839 | 4,027 | 4,646 | 3,220 |
| Bauchi | 20/20 | 100 | 2,368 | 2,590 | 2,771 | 2,187 |
| Bayelsa | 8/8 | 40 | 967 | 976 | 1,102 | 841 |
| Benue** | 14/23 | 70 | 1,561 | 1,891 | 2,181 | 1,271 |
| Cross River | 18/18 | 87 | 2,399 | 2,544 | 3,247 | 1,696 |
| Ekiti | 16/16 | 75 | 1,658 | 1,865 | 2,144 | 1,379 |
| FCT** | 4/6 | 20 | 505 | 498 | 520 | 483 |
| Jigawa | 27/27 | 135 | 2,954 | 3,575 | 3,015 | 3,514 |
| Kaduna | 23/23 | 115 | 2,742 | 3,119 | 2,987 | 2,874 |
| Kano | 44/44 | 220 | 5,106 | 5,898 | 5,408 | 5,596 |
| Katsina | 34/34 | 170 | 3,901 | 4,435 | 3,531 | 4,805 |
| Kebbi** | 20/21 | 100 | 1,505 | 3,349 | 2,701 | 2,153 |
| Kogi | 21/21 | 105 | 2,541 | 2,731 | 2,872 | 2,400 |
| Lagos** | 19/20 | 95 | 2,437 | 2,337 | 1,712 | 3,062 |
| Niger | 25/25 | 132 | 3,143 | 4,054 | 4,503 | 2,694 |
| Osun | 30/30 | 149 | 3,703 | 3,876 | 4,737 | 2,842 |
| Oyo | 33/33 | 165 | 3,983 | 4,127 | 4,488 | 3,622 |
| Rivers | 23/23 | 115 | 2,887 | 2,833 | 2,399 | 3,321 |
| Taraba** | 8/16 | 37 | 892 | 955 | 871 | 976 |
| Yobe* | 15/17 | 75 | 1,711 | 1,990 | 1,764 | 1,937 |
| Total | 433/456 | 2,160 | 50,802 | 57,670 | 57,599 | 50,873 |

** States where some LGAs were previously mapped

* State where some LGAs were not mapped due to insecurity

3.2 Parasitological Findings

Out of the 108,472 pupils sampled, 10,349 (9.5%) were infected with schistosomes, 29,269 (27.0%) with STHs and 2,163 (2.0%) were co-infected with schistosomes and STHs. In the 19 States and FCT surveyed for schistosomiasis, the highest prevalence occurred in Niger State (26.1%), followed by Kebbi State (21.9%) and FCT (20.3%). Lowest values were recorded in Rivers and Akwa Ibom States with prevalence of 0.1% and 0.3% respectively. The prevalence of STHs was highest in Akwa Ibom State (58.4%) followed by Oyo (47.2%) and Osun (45.2%) and lowest in Yobe (1.4%) and Taraba (5.6%) States. Of the infected, 2,163 (1.99%) co-infected, Niger State had the highest prevalence of 8.96% followed by FCT 4.19%. Rivers State had the lowest prevalence of 0.03% (Table 3 and figure 3).

Table 3: Prevalence of Schistosomiasis, Soil Transmitted Helminths and Co-Infection

| State | No of persons examined | Schistosomiasis | | STH | | Schistosomiasis and STH co-infection |
|--------------|------------------------|---------------------|----------------------|----------------------|------------------------|--------------------------------------|
| | | No.(%) infected | 95% CI | No.(%) infected | 95% CI | No.(%) infected |
| Akwa-Ibom | 7,866 | 22 (0.3) | (0.18 – 0.43) | 4,590 (58.4) | (57.25 – 59.44) | 12(0.15) |
| Bauchi | 4,958 | 675 (13.6) | (12.67 – 14.6) | 496 (10.0) | (9.19 – 10.88) | 58 (1.17) |
| Bayelsa | 1,943 | 17 (0.9) | (0.52 – 1.42) | 639 (32.9) | (30.81 – 35.04) | 10 (0.51) |
| Benue | 3,452 | 451 (13.1) | (11.96 – 14.24) | 872 (25.3) | (23.82 – 26.75) | 135 (3.91) |
| Cross River | 4,943 | 283 (5.7) | (5.11 – 6.42) | 1,209 (24.5) | (23.27 – 25.69) | 45 (0.91) |
| Ekiti | 3,523 | 8 (0.2) | (0.11 – 0.47) | 1,084 (30.8) | (29.25 – 32.33) | 4 (11) |
| FCT | 1,003 | 204 (20.3) | (17.92 – 22.99) | 193 (19.2) | (16.87 – 21.85) | 42 (4.19) |
| Jigawa | 6,529 | 743 (11.4) | (10.62 – 12.18) | 404 (6.2) | (05.62 – 06.81) | 66 (1.01) |
| Kaduna | 5,861 | 811 (13.8) | (12.97 – 14.76) | 1,279 (21.8) | (20.77 – 22.90) | 122 (2.08) |
| Kano | 11,004 | 1,531 (13.9) | (13.27 – 14.57) | 1,923 (17.5) | (16.78 – 18.21) | 307 (2.79) |
| Katsina | 8,336 | 944 (11.3) | (10.65 – 12.02) | 872 (10.5) | (9.82 – 11.14) | 127 (1.52) |
| Kebbi | 4,854 | 1,062 (21.9) | (20.73 – 23.08) | 480 (9.9) | (9.07 – 10.77) | 85 (1.75) |
| Kogi | 5,272 | 149 (2.8) | (2.41 – 3.32) | 1,481 (28.1) | (26.88 – 29.33) | 51 (0.97) |
| Lagos | 4,774 | 41 (0.9) | (0.63 – 1.18) | 1,341 (28.1) | (26.82 – 29.39) | 12 (0.25) |
| Niger | 7,197 | 1,879 (26.1) | (25.1 – 27.14) | 2,531 (35.2) | (34.07 – 36.29) | 645 (8.96) |
| Osun | 7,579 | 405 (5.3) | (4.88 – 5.88) | 3,426 (45.2) | (44.08 – 46.33) | 203 (2.68) |
| Oyo | 8,110 | 435 (5.4) | (4.88 – 5.88) | 3,828 (47.2) | (46.11 – 48.29) | 216 (2.66) |
| Rivers | 5,720 | 7 (0.1) | (0.05 – 0.26) | 2,467 (43.1) | (41.84 – 44.43) | 2 (0.03) |
| Taraba | 1,847 | 103 (5.6) | (4.6 – 6.75) | 103 (5.6) | (4.60 – 6.75) | 12 (0.65) |
| Yobe | 3,701 | 579 (15.6) | (14.49 – 16.86) | 51 (1.4) | (1.04 – 1.82) | 9 (0.24) |
| Total | 108,472 | 10,349 (9.5) | (9.37 – 9.72) | 29,269 (27.0) | (26.72 – 27.25) | 2,163 (1.99) |




Figure 3: Schistosomiasis and STHs endemic areas

In the LGAs of each State, it was also found that the range of prevalence of each disease varied (Table 4). With respect to schistosomiasis, LGAs prevalence in four States were in the low risk range, LGAs in 13 States ranged from low to moderate risk while in two States; Niger and Kebbi, the prevalence ranged from low to high as stated in WHO treatment guidelines for schistosomiasis and STHs (Appendix 6). STHs disease prevalence range in the LGAs of the 19 States and FCT surveyed showed wider variation. LGAs in four States were in the clinical case management. In seven States, the LGA prevalence rate ranged from clinical to low risk levels. Only Benue State had all its LGAs at low risk range. Five States had clinical to high risk prevalence levels in the LGAs while two States; Akwa Ibom and Osun had prevalence levels ranging between low and high. Figures 4 and 5 show the prevalence of schistosomiasis and STHs respectively.

Table 4: Range of Schistosomiasis and STHs Prevalence by State

| State | No of LGAs Surveyed | Prevalence Range (%) | |
|-------------|---------------------|----------------------|-------------|
| | | Schistosomiasis | STHs |
| Akwa Ibom | 31 | 0.0 - 1.2 | 25.1 - 91.4 |
| Bauchi | 20 | 0.0 - 33.9 | 1.6 - 19.3 |
| Bayelsa | 8 | 0.0 - 3.4 | 14.9 - 59.2 |
| Benue | 14 | 1.4 - 24.8 | 20.2 - 36.8 |
| Cross River | 18 | 0 - 32.8 | 12.5 - 50.2 |
| Ekiti | 16 | 0.0 - 2.5 | 15.8 - 48.9 |
| FCT | 4 | 7.5 - 27.9 | 16.1 - 25.5 |
| Jigawa | 27 | 0.5 - 31.7 | 0 - 15.4 |
| Kaduna | 23 | 1.5 - 44.2 | 4.2 - 40.0 |
| Kano | 44 | 1.2 - 39.0 | 5.6 - 34.0 |
| Katsina | 34 | 0 - 28.9 | 0.8 - 40.6 |
| Kebbi | 20 | 0.8 - 68.3 | 3.8 - 22.0 |
| Kogi | 21 | 0 - 21.4 | 16.2 - 39.0 |
| Lagos | 19 | 0 - 3.2 | 0 - 44.8 |
| Niger | 25 | 2.8 - 51.6 | 13.7 - 64.7 |
| Osun | 30 | 0 - 16.2 | 24 - 71.1 |
| Oyo | 33 | 0 - 19.6 | 16.7 - 67.7 |
| Rivers | 23 | 0 - 0.8 | 17.6 - 85.9 |
| Taraba | 8 | 1.0 - 11.1 | 1.6 - 12.9 |
| Yobe | 15 | 1.2 - 39.6 | 0 - 3.3 |




Figure 4: Prevalence of Schistosomiasis by LGA




Figure 5: Prevalence of STHs by LGA

3.1.1 Schistosomiasis Result

Schistosoma haematobium (82%) was the predominant species in the survey compared to *Schistosoma mansoni* (18%) as shown in Figure 6. Of the total pupils examined, 8.1% were positive for *S. haematobium* with highest prevalence occurring in Kebbi (21.7%) followed by Niger (19.6%) and Yobe (15.6%) States. For *S. mansoni*, 1.8% pupils were positive, FCT and Niger States ranked highest with prevalence of 10.5% and 9.4% respectively while Yobe State had no *S. mansoni*. Akwa Ibom, Lagos, Katsina and Rivers States had 0.1% prevalence each (Figures 6, 7 and 8).




Figure 6: Proportion of *S. haematobium* and *S. mansoni* in the study area




Figure 7: Prevalence of *S. haematobium* and *S.mansoni* by State




Figure 8: Schistosomiasis Point Prevalence maps

3.1.2 Soil Transmitted Helminths (STHs) Results

All three STHs species *Ascaris lumbricoides*, Hookworm and *Trichuris trichiura* were observed in the present survey. The most prevalent species was hookworm (47%), *A. lumbricoides* (42%) and *T. trichiura* (11%). All the 19 States and FCT surveyed showed presence of *A. lumbricoides* and Hookworm infections. Only Yobe State had no prevalence of *T. trichiura*. Osun State followed by Oyo and Akwa Ibom States had the highest prevalence of *A. lumbricoides* with 40.9%, 35.9% and 31.9% respectively. Hookworm infection was highest in Akwa Ibom State with 40.3% followed by Niger State with 32.6% and the lowest in Yobe State with 0.7%. Rivers State recorded highest prevalence in *Trichuris* infection with 19.1% and followed closely by neighbouring Bayelsa State with 15.5% (Figure 9, 10 and 11). Of the 4.0% that had multiple STHs infection, Akwa Ibom State had highest prevalence of 20.8%. The lowest prevalence of 0.04% was recorded in Kebbi State.




Figure 9: Proportion of the three species of STHs in the Study Area




Figure 10: Prevalence of *A. lumbricoides*, Hookworm and *T. trichiura*




Figure 11: STHs Point Prevalence maps

3.1.3 Prevalence of schistosomiasis and STH by Sex

Of the total pupils infected with schistosomiasis, 65% were males and 35% females. Niger State had highest prevalence of infection both in males (29.4%) and females (21.9%). Rivers State had the least infection in both sexes (Figure 12 and 13). There is a statistical significant association with schistosomiasis infection by sex ($\chi^2 = 957.37, P < 0.05$).




Figure 12: Proportion of pupils infected with Schistosomiasis by sex




Figure 13: Prevalence of Schistosomiasis by sex

For STHs, of the total infected pupils, 53% were males and 47% females. Akwa Ibom State had highest prevalence in both males (59.8%) and females (56.8%) while Yobe State had the least prevalence for both sexes 1.2% for males and 1.6% females (Figures 14 & 15). There is a statistically significant association between sex and STHs infection ($\chi^2 = 101.49, P < 0.05$).




Figure 14: Proportion of Pupils infected with STHs by sex




Figure 15: Prevalence of STHs by sex

3.1.4 Prevalence of schistosomiasis and STH by Age group

Schistosomiasis was found to be more prevalent among the 11-16 years age group (55%) as compared to the 5-10 years age group (45%) as shown in figure 16. In both age groups examined, Niger State had the highest prevalence of schistosomiasis with 24.1% and 29.4% for 5-10 and 11-16 years respectively. The lowest prevalence of 0.04% and 0.2% were recorded in Rivers State for both age groups respectively (Figure 17). Statistically, it was also observed that there was significant difference between prevalence of disease of the age groups ($\chi^2 = 139.48, P < 0.05$).




Figure 16: Proportion of schistosomiasis infected pupils by age group




Figure 17: Prevalence of schistosomiasis by age group

STHs were found to be more prevalent among the 5-10 years age group (51%) as compared to the 11-16 years age group (49%). Akwa Ibom had the highest prevalence for both age groups; 56.2% for 5-10 years and 61.5% for 11-16 years. The lowest prevalence was found in Yobe State with 1.4% and 1.3% for both age groups respectively (Figures 18 and 19). There was statistically significant difference between the age groups ($\chi^2 = 11.12, P < 0.05$).




Figure 18: Proportion of STHs infected pupils by age group




Figure 19: Prevalence of STHs infected pupils by age group

3.1.5 Intensity of Infection

The intensity levels (expressed as numbers of eggs per ml of urine or gram of faeces) of the various species of parasites recorded in this mapping survey are shown in table 5. The result shows that there were generally more cases of light and moderate intensities of infection. Heavy intensity levels were nearly equal for *S. haematobium* (24.31%) and *S. mansoni* (23.48%). However, heavy intensity of *S. haematobium* was higher in males (25.02%) than females (22.64%). Conversely, females (25.41%) recorded cases of heavy intensity of *S. mansoni* than males (22.01%). The intensity levels of *S. haematobium* showed statistical significant difference ($P < 0.05$) with respect to sex in this survey.

For the STHs, hookworm (2.94%) recorded more cases of heavy infection followed by *A. lumbricoides* (1.40%) and *T. trichuria* (0.50%) had the least. The intensity levels of *A. lumbricoides* showed statistical significant difference ($P < 0.05$) with respect to sex in this survey.

Table 5: Parasite intensity in eggs per gram (epg) of faeces or per 10 ml of urine

| Parasite species | Level of intensity | Number (%) of intensity of infection | | |
|------------------------|---------------------------------------|--------------------------------------|---------------|----------------|
| | | Male | Female | Total |
| <i>S. haematobium</i> | Light Infection (<50 eggs/ml) | 4,605 (74.98) | 2,033 (77.36) | 6,638 (75.69) |
| | Heavy Infection (50 eggs/ml) | 1,537 (25.02) | 595 (22.64) | 2,132 (24.31) |
| <i>S. mansoni</i> | Light Infection (1 – 99 epg) | 461 (41.09) | 325 (38.42) | 786 (39.94) |
| | Moderate Infection (100-399 epg) | 414 (36.90) | 306 (36.17) | 720 (36.59) |
| | Heavy Infection (>400 epg) | 247 (22.01) | 215 (25.41) | 462 (23.48) |
| <i>A. lumbricoides</i> | Light Infection (1-4,999 epg) | 5,721 (76.28) | 5,061 (74.42) | 10,782(75.39) |
| | Moderate Infection (5,000-49,999 epg) | 1,681 (22.41) | 1,638 (24.08) | 3,319 (23.21) |
| | Heavy Infection (>50,000 epg) | 98 (1.31) | 102 (1.50) | 200 (1.40) |
| Hookworm | Light Infection (1-1,999 epg) | 8,883 (92.70) | 6,143 (92.03) | 15,026 (92.42) |
| | Moderate Infection (2,000-3,999 epg) | 443 (4.62) | 311 (4.66) | 754 (4.64) |
| | Heavy Infection (>4,000 epg) | 257 (2.68) | 221 (3.31) | 478 (2.94) |
| <i>T. trichiura</i> | Light Infection (1-999 epg) | 1,684 (88.45) | 1,484 (86.43) | 3,168 (87.49) |
| | Moderate Infection (1,000-9,999 epg) | 212 (11.13) | 223 (12.99) | 435 (12.01) |
| | Heavy Infection (>10,000 epg) | 8 (0.42) | 10 (0.58) | 18 (0.50) |

3.2 Implication of Findings on Intervention

Results of the survey showed that of 433 LGAs mapped, 359 (83%) required interventions for schistosomiasis (Appendix 7). Furthermore, in eight States and FCT, all the LGAs mapped (180) require treatment. Rivers State has the least number of LGAs (26%) requiring treatment. A total of 202 LGAs fall within the low risk category, 153 moderate and four LGAs were high risk category. The four LGAs in the high risk category are found in Niger (2) and Kebbi (2) States (Table 6 and Figure 20).

Table 6: LGAs requiring Intervention for Schistosomiasis

| State | No of LGAs mapped | Number of LGAs requiring Intervention | | | |
|--------------------|-------------------|---------------------------------------|----------------------------|----------------|--|
| | | Low Risk (<10%) | Moderate Risk (10 - 49.9%) | High Risk ≥50% | Total number (%) of LGA requiring intervention |
| Akwa Ibom | 31 | 15 | 0 | 0 | 15 (48) |
| Bauchi | 20 | 8 | 11 | 0 | 19 (95) |
| Bayelsa | 8 | 6 | 0 | 0 | 6 (75) |
| Benue | 14 | 5 | 9 | 0 | 14 (100) |
| Cross-River | 18 | 6 | 5 | 0 | 11 (61) |
| Ekiti* | 16 | 3 | 0 | 0 | 3 (19) |
| FCT | 4 | 1 | 3 | 0 | 4 (100) |
| Jigawa | 27 | 14 | 13 | 0 | 27 (100) |
| Kaduna | 23 | 9 | 14 | 0 | 23 (100) |
| Kano | 44 | 19 | 25 | 0 | 44 (100) |
| Katsina | 34 | 13 | 18 | 0 | 31 (91) |
| Kebbi | 20 | 5 | 13 | 2 | 20 (100) |
| Kogi | 21 | 13 | 3 | 0 | 16 (76) |
| Lagos | 19 | 13 | 0 | 0 | 13 (68) |
| Osun | 30 | 24 | 4 | 0 | 28 (93) |
| Oyo | 33 | 27 | 4 | 0 | 31 (94) |
| Rivers | 23 | 6 | 0 | 0 | 6 (26) |
| Taraba | 8 | 7 | 1 | 0 | 8 (100) |
| Yobe | 15 | 6 | 9 | 0 | 15 (100) |
| Niger | 25 | 2 | 21 | 2 | 25 (100) |
| Total | 433 | 202 | 153 | 4 | 359 (83) |

* Ekiti was not mapped for *S. haematobium* in this survey




Figure 20: Intervention planning map for Schistosomiasis in 19 States and FCT.

A total of 237(55%) LGAs require some form of intervention for STHs as shown in Table 7 and Figure 21. All the LGAs mapped in Akwa Ibom, Benue and Osun States (75 LGAs) qualify for intervention. However, no LGA was found to qualify for intervention in Bauchi, Jigawa, Taraba and Yobe States. Case-based management is required in 191 LGAs of the surveyed States, 177 LGAs fall within the low risk and 60 LGAs in the high risk categories (Appendix 7).

Table 7: LGAs requiring Intervention for Soil Transmitted Helminths (STHs)

| State | No of LGAs mapped | Number of LGAs requiring Intervention | | | |
|--------------|-------------------|---------------------------------------|-----------------------|------------------|---|
| | | Case-based Mgt (<20%) | Low Risk (20 - 49.9%) | High Risk (≥50%) | Total No. (%) of LGA requiring intervention |
| Akwa Ibom | 31 | 0 | 9 | 22 | 31 (100) |
| Bauchi | 20 | 20 | 0 | 0 | 0 (0) |
| Bayelsa | 8 | 3 | 3 | 2 | 5 (63) |
| Benue | 14 | 0 | 14 | 0 | 14 (100) |
| Cross-River | 18 | 9 | 8 | 1 | 9 (50) |
| Ekiti | 16 | 1 | 15 | 0 | 15 (94) |
| FCT | 4 | 3 | 1 | 0 | 1 (25) |
| Jigawa | 27 | 26 | 0 | 0 | 0 (0) |
| Kaduna | 23 | 9 | 14 | 0 | 14 (61) |
| Kano | 44 | 27 | 17 | 0 | 17 (39) |
| Katsina | 34 | 30 | 4 | 0 | 4 (12) |
| Kebbi | 20 | 18 | 2 | 0 | 2 (10) |
| Kogi | 21 | 3 | 18 | 0 | 18 (86) |
| Lagos | 19 | 15 | 3 | 0 | 3 (16) |
| Osun | 30 | 0 | 20 | 10 | 30 (100) |
| Oyo | 33 | 1 | 19 | 13 | 32 (97) |
| Rivers | 23 | 2 | 11 | 9 | 20 (87) |
| Taraba | 8 | 8 | 0 | 0 | 0 (0) |
| Yobe | 15 | 13 | 0 | 0 | 0 (0) |
| Niger | 25 | 3 | 19 | 3 | 22 (88) |
| Total | 433 | 191 | 177 | 60 | 237 (55) |




Figure 21: Intervention planning map for STHs in 19 States and the FCT

3.3 Defecation practices and water contact activities

Table 8 shows prevalence of schistosomiasis and STHs in relation to place of defecation by pupils surveyed. The highest prevalence for schistosomiasis and STHs was among respondents who defecated around the school compound (11.1% and 31.0% respectively). The second highest prevalence occurred among those who defecate outside school environment (10.7% for schistosomiasis and 26.5% for STHs). The State level prevalence is as shown on Appendix 8.

Table 8: Prevalence of schistosomiasis and STHs in relation to places of defecation.

| Defecation Practices | No of Persons Examined | Schistosomiasis | STHs |
|----------------------------|------------------------|----------------------|-----------------------|
| | | No (%) Infected | No (%) Infected |
| In the School Toilet | 45,726 | 3,501 (7.66) | 11,314 (24.74) |
| Around school Compound | 29,847 | 3,319 (11.12) | 9,250 (30.99) |
| Outside School Environment | 32,899 | 3,529 (10.73) | 8,705 (26.46) |
| Total | 108,472 | 10,349 (9.54) | 29,269 (26.98) |

Table 9 and Appendix 9 showed that all respondents had some form of contact with water. The activities include bathing, fishing, washing, swimming, playing, fetching water, crossing water and others. The prevalence of schistosomiasis was highest among those who perform swimming activity

(17.0%) followed by those who play in water (11.3%). For STHs the highest prevalence was among those who bath (34.3%) and those who fish (33.9%).

Table 9: Prevalence of Schistosomiasis in relation to water contact activities

| Water Contact Activities | No of Respondents | Schistosomiasis | STHs |
|--------------------------|-------------------|----------------------------|----------------------------|
| | | No (%) of Persons Infected | No (%) of Persons Infected |
| Bathing | 13,835 | 1,351(9.8) | 4,739 (34.3) |
| Washing | 20,689 | 2,106 (10.2) | 6,160 (29.8) |
| Fishing | 3,535 | 349 (9.9) | 1,199 (33.9) |
| Crossing Water | 3,253 | 279 (8.6) | 917 (28.2) |
| Fetching Water | 14,804 | 1,124 (7.6) | 4,481 (30.3) |
| Playing | 14,364 | 1,624 (11.3) | 3,223 (22.4) |
| Swimming | 6,590 | 1,119 (17.0) | 1,970 (29.9) |
| Others | 31,402 | 2,397 (7.6) | 6,580 (21.0) |
| Total | 108,472 | 10,349 (9.5) | 29,269 (27.0) |

4.0 DISCUSSIONS

4.1 Discussion of Results

The results of the Schistosomiasis and STHs epidemiological mapping in the 19 States and FCT provided insight to the disease distribution and intensity by age group and sex for pupils in the surveyed area.

There are 456 LGAs in the 19 States but 21 LGAs (in Benue, FCT, Lagos, Kebbi and Taraba) had previously been mapped by government and partners. The intention of this survey was to complete the mapping in the remaining 435 LGAs; however, two LGAs in Yobe State could not be mapped due to insecurity.

The findings on schistosomiasis revealed that the overall prevalence was within the low risk range (9.5%), although data captured by LGA showed high levels of prevalence in some LGAs within some States. High salinity waters may not support the viability of freshwater snail hosts of schistosomes and may account for the low prevalence observed in Rivers, Akwa Ibom, Bayelsa and Lagos States that border the Atlantic and have brackish water bodies. The finding of this present survey agrees with other studies by Osama, 2009; Leveque *et al.*, 1978; Kefford and Nugegoda (2005). However, in Cross River also a coastal State, five LGAs had moderate risk. These LGAs are situated in the northern part of the State bordering Benue State where the infection was also moderate. The low prevalence in Ekiti State could be explained by the deworming programme for schistosomiasis which was launched in 2010 although treatment has not been consistent over the years.

Regarding STHs, the overall prevalence rate was low, but LGA rates in some State were greater than 50% which is WHO classification for high risk (WHO, 2011). Most of the LGAs with high risk were found in States in the southern part of the country except Niger State in the North Central zone. The survey showed higher prevalence of STHs than schistosomiasis which was in contrast to findings in similar studies in Zimbabwe where schistosomiasis was 22.7% and STHs 5.5% (Midzi *et al.*, 2014)

Prevalence of infection shows the parasites burden in the population while intensity estimates the worm burden per host. The ability of a parasite to elicit disease symptoms depends on its intensity in the host and a combination of other factors such as the physiological state of the host and the presence of other parasites. Although cases of heavy intensity were few in this survey, those of moderate and light infections were quite high and could graduate into heavy intensity if they remain untreated. Studies have demonstrated a relationship between the initial worm burden and growth after treatment (Mohammed *et al.*, 2002). Children with higher worm burden were observed to gain less weight than those with lower worm burden.

Co-infection of schistosomiasis and STHs was observed in all the States sampled except in Rivers State. The range of the prevalence of schistosomiasis and STHs in the LGAs of the States was in agreement with the findings from other studies that schistosomiasis is a focal disease even in one State where its

prevalence varied from LGA to LGA (FMOH 2013). Conversely, STH are ubiquitous and the low prevalence range in Jigawa and Yobe States could be attributed to on-going annual Lymphatic Filariasis treatment with Albendazole in some parts of the State in the last five years. The range provided information on the LGAs requiring intervention as well as the type of treatment required.

Schistosoma haematobium was more prevalent than *S. mansoni* in the present survey and agrees with similar studies in Nigeria. Some States such as Kebbi, Niger and Yobe with very high prevalence of *S. haematobium* may be linked to presence of freshwater bodies and dams dotting such areas. These dams and other natural water bodies support irrigation and provide recreational facilities for school age children who frequent these sources for different activities that can lead to exposure to infection. The water bodies often are the only source of water for domestic uses and also serve as social centres for the communities. Dams and other freshwater bodies have been implicated in epidemiology of schistosomiasis in Nigeria and Africa (Steinmann *et al.*, 2006). Nigeria is still endemic for schistosomiasis and scale-up provision of infrastructure especially potable water may reduce contact with infected waters and further support the elimination programme of the government.

Ascaris, hookworms and *Trichuris* are prevalent in Nigeria and associated with poor infrastructure and unsanitary habits. The presence of multiple infections with these STHs in school age children has been observed in similar studies globally including Nigeria (Worrell *et al.*, 2013, Ojurongbe *et al.*, 2014). These impose high burden on the infected pupils causing chronic morbidity, cognitive impairment and school absenteeism (Lobato *et al.*, 2012). The high prevalence of *Ascaris* and hookworms observed in Osun, Oyo, Akwa-Ibom, Bayelsa and other rain forest zone States may be attributed to favourable climatic conditions of adequate moisture and relative humidity and poor sanitary condition which enhance transmission. In contrast, a State like Yobe in northern savannah zone where there is low moisture content had very low prevalence.

Males were significantly more susceptible to schistosome infections than females as observed in this survey. This has been linked to increased and prolonged water contact activities by males such as swimming, fishing and bathing especially during the peak hours of cercariae shedding by snail hosts which are risk factors in exposure to infection. The females are engaged in lower risk activities of washing and fetching water which usually occur in the early mornings and late evenings when cercariae shedding is low (Anya & Okoronkwo 1991). Moreover there is limited exposure of the body during these activities.

The high prevalence of schistosomiasis among pupils 11 – 16 years old is suggestive of possible frequent water contact activities in this age group. Pupils of this age range frequent water bodies for recreational activities like swimming than those in the younger ages (Worrell, *et al.* 2013). This is exemplified by the fact that in Niger State where the prevalence of schistosomiasis was high, it

occurred more in this age group (11 – 16 years), particularly, along the major water points in the State. Age-related activities and behaviour, frequency of exposure and development of immunity are known to play important role in the distribution of schistosomiasis and STHs infections (Daniel *et al.*, 2007). Conversely, higher prevalence of STHs was recorded in younger age group (5 - 10 years). The statistical significance with respect to age group is a clear indication that pupils within this age group are more exposed to the sources of infection than the older ones. High infection rates with STHs have been attributed to poor sanitary conditions and hygiene, low nutritional status (Otubanjo and Mafe, 2002), climatic conditions such as temperatures and soil factors such as salinity (Bosompem, 2004). A proposed causal link between hygiene and intestinal parasitic infections (Allison and Larson 2002) features water contact-base activities and direct consumption of contaminated water as major sources of infections.

4.2 Implications of Findings

The implication of the findings of this survey for treatment intervention for schistosomiasis at the LGAs' level is presented at three prevalence categories (WHO, 2011): high-risk areas (where prevalence of infection in school-age children is $\geq 50\%$), moderate-risk areas (where prevalence of infection in school-age children is $>10\%$ but $<50\%$) and low-risk areas (where prevalence of infection in school-age children is $<10\%$). The estimated population at risk in the 19 States and FCT is 24,195,603 persons for schistosomiasis.

This categorization of treatment intervention is required for planning purposes. The level of prevalence of infection in an area is used to determine the frequency of treatment. By this, four LGAs in this survey required annual treatment of all school-age children and adults considered to be at risk. One hundred and fifty-three LGAs require treatment once every two years for all school age children and adults considered to be at risk. Two hundred and two LGAs are qualified for two treatment intervention during their primary schooling years. An estimated 21,257,368 persons are at risk of STHs infections in the 19 States and FCT.

WHO (2011) however, presents only two categories for the STHs (high-risk and low-risk areas). Sixty LGAs fall within the high-risk areas while 177 LGAs were within the low-risk areas. The implication is twice yearly treatment for school age children in these 60 LGAs. In addition, treatment should also be administered to other at-risk individuals including pre-school age children, women of child-bearing age and adults in high risk occupation. This shows the magnitude of treatment intervention required for control and elimination purposes.

In addition to preventive chemotherapy, interventions should include the implementation of other components of the PHASE strategy. The findings of low numbers of LGAs requiring intervention in

Rivers State may be partly due to the recent scale up/development of schools infrastructure and improvement in sanitary conditions of the schools.

The places of defaecation were categorised into three; in the school toilet, around school compound and outside school environment. The results showed that pupils using any of the three places for defaecation were infected with either or both parasites. However, the prevalence was lowest among those pupils who defaecated in the school toilet. Adebote *et al.*, (2005), reported high prevalence of helminth ova in faeces deposited at different locations near human habitations and concluded that poor personal hygiene and environmental sanitation, which can occur as a consequence of indiscriminate defaecation and the use of human excreta as fertilizer on farms and vegetable gardens (Umoh *et al.* 2001), may result in spread of helminth infections.

4.3 Some Key Achievements of the Project

- The CIFF investment supported the strengthening of the office of the National Coordinator for Schistosomiasis and STHs Control Programme. This has ensured a well-equipped and functional office for proper coordination of the programme in Nigeria.
- The capacity of health workers was developed/strengthened across the tiers of Government to respond to NTDs programme direction in the country. It also created opportunity to showcase multi-sectoral approach and collaboration for result.
- The support was able to leverage additional funding (\$80,000) from RTI/ENVISION to map additional LGAs in 2 States (Cross Rivers and Ondo) where mapping had not been completed.
- Sightsavers also funded (\$25,715) for the completion of the mapping of schistosomiasis and STHs in Benue State.
- There was also the mapping of other States (Bauchi, Taraba and Niger) with DFID funding from the DFID/GTMP.
- The investment has also helped Nigeria to realign its control/elimination road map with the WHO recommended timeline for NTDs.

5.0. CHALLENGES/CONSTRAINTS

There were some challenges encountered during the mapping project. They have been grouped into five categories as follows:

5.1 Logistics

- Lack of counterpart funding from the State which led to inadequate mobilization and sensitization of the schools and communities.
- Inefficient logistics including project vehicles in some States for project monitoring and implementation.
- Distances between selected schools were sometimes far hence the teams arrived their second schools late. This was worse if the two schools are in the riverine areas.
- Unanticipated fuel scarcity and high cost of fuel in some of the States impacted on overall project schedule and cost.

5.2 Mobilisation/Misconception

Inadequate mobilisation of some of the host communities coupled with misconceptions in some areas about handling of human wastes led to delays and in some cases rejection by respondent to participate. Some team members were also harassed and manhandled by communities who questioned the implication of taking their children's specimen away from the village.

5.3 Communication/Technology Challenges

- Challenges of communication reduced the level of communication between the supervisors/consultants and the teams during the field work. It was very important to track the team leaders and have real-time information on the progress of work.
- Difficulties with uploading of information in the phone were experienced. In some areas, other internet service providers had comparative advantage over MTN which was solely used by the programme.

5.4 Access challenges to schools in some LGAs

The Terrain in some LGAs made the schools difficult to reach and a team ends up visiting only one school in a day and not two as in the plan. Some teams had to cross the ocean to reach the riverine areas and mapping took up to two days per school in some cases. Also, some children did not resume promptly from the first term school vacation which had just ended before the commencement of the mapping exercise causing delays in completion of the mapping exercise in some States.

5.5 Security Challenges

Insecurity in some parts of the country and the concomitant nationwide political campaigns during the period of this exercise were reasons for some of the delays in securing approvals to commence the project in some States. Residents were displaced in some LGAs and the school children were not available for the survey in other areas. These unforeseen situations hindered timely achievement of set targets of the mapping project.

6.0. RECOMMENDATIONS

Having successfully conducted the mapping of schistosomiasis and STHs in 19 States and FCT, there is ample evidence for appropriate intervention. It is recommended that the FMOH should carry out the following:

1. Disseminate the results of the mapping survey at all levels including WHO and other global NTDs medicines donors using appropriate channels
2. Strengthen collaboration with relevant sectors and partners (FMOH, SMOH, SUBEB and other Partners) to commence the implementation of PHASE in all endemic LGAs
3. Establish electronic platform at the FMOH office for data transmission, storage and analysis with respect to future schistosomiasis and STH interventions.
4. Enhanced Advocacy, mobilization and multi-sectoral approach to ensure maximum community participation.
5. Leverage funding support from Government at all levels, NGOs and corporate organisations using the results of the mapping.
6. Provide field operational vehicles at the national office for supportive supervision and monitoring of project implementation.

7.0 CONCLUSION

The mapping of schistosomiasis and STHs in 19 States and the Federal Capital Territory of Nigeria was successfully completed and has achieved all the set objectives. Data generated from this crucial survey provided vital evidence for appropriate and sustainable intervention by Government in collaboration with our highly esteemed NTDs partners. It is the right of every child in Nigeria to enjoy good health and the time to intervene is now.


REFERENCES

1. Adebote, D.A., Oniye, S.J. and Aliyu, I.S. (2005): Occurrence of helminth ova in peridomestic human excreta in Zaria, Northern Nigeria. *The Zoologist*, 3: 8-14.
2. Allison, E.A. and Larson, E. (2002): What is the evidence for a causal link between hygiene and infections? *Lancet Infectious Diseases*, 2: 103-110.
3. Anya and Okoronkwo (1991): Observation on the optimal conditions for infection and development of *Schistosoma haematobium* (Trematoda, Schistosomatides) in the snail host in Eastern Nigeria, 1991.
4. Bosompem, K.M. (2004): Infants schistosomiasis in Ghana: a survey in an irrigation community. *Tropical Medicine and International Health*, 9: 917-922.
5. Daniel, E.E., Robert, W.S. and Joel, W. (2007): Helminths as governors of immune-mediated inflammation. *International Journal for Parasitology*, 37: 457-464.
6. Feachem, R.G., Guy, W.M., Harrison, S., Iwago, K.O., Marshall, T., Mbere, N., Muller, R. and Wright, A.M. (1983): Excreta disposal facilities and intestinal parasitism in urban Africa: Preliminary studies in Botswana, Ghana and Zambia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 77: 515-521.
7. FMOH (2013-2017): Nigeria Master Plan for Neglected Tropical Diseases: Federal Ministry of Health, Abuja.
8. FMOH (2013): National Protocol for Epidemiological Mapping and baseline survey of Schistosomiasis and Soil Transmitted Helminths in Nigeria. 55pp.
9. Hotez, P.J., Herricks, J.R. (2015): Helminth Elimination in the Pursuit of Sustainable Development Goals: A "Worm Index" for Human Development. *PLoS Negl Trop Dis* 9(4): e0003618. doi:10.1371/journal.pntd.0003618
10. Hotez P.J, Aruna K. (2009): Neglected Tropical Diseases in Sub-Sahara Africa: Review of their Prevalence, Distribution and Disease Burden. *Plos NTDs* 3(8) doi: 10.1371/Journal.pntd.0000412.
11. Lengeler, C., Mshinda, H., Morona, D. & Desavigny, D. (1993): Urinary schistosomiasis: testing with urine filtration and reagent sticks for haematuria provides a comparable prevalence estimate. *Acta Trop*, 53, 39-50.
12. Lobato L., Miranda, A., Faria, I.M., Bethony, J.M., Gazzinelli, M.F. (2012): Development of Cognitive abilities of children infected with helminths through health education. *Revista da Sociedade Brasileira de Medicina Tropical* 45 (4): 514-519, July – August 2012.
13. Midzi N., Mduluma, T., Charimari, L., Mhlanga, G., Chimbari, M.J., Tshuma, C., Manangazira, P., Munyati, S.N., Phiri, I., Mutambu, S.L., Midzi, S.S., Ncube, A., Muranzi L.P.,

- Rusakaniko, S., Mutapi, F. (2014): Distribution of Schistosomiasis and Soil Transmitted helminthiasis in Zimbabwe: Towards a National Plan of Action for Control and Elimination. *PLoS Neglected Tropical Diseases* 8(8): e3014. Doi: 10.1371/journal.pntd.0003014.
14. Mohammed, F., Akbar, M. Jamalain, F. and Ahmad, G. (2002): Evaluation of two years mass chemotherapy against ascariasis in Hamadan, Islamic Republic of Iran. *International Journal of Public Health*, 80(5): 399-402
 15. National Population Commission (NPC 2006): The Nigeria Census Population.
 16. Ojurongbe O., Oyesiji, K.F., Ojo, J.A., Adewale, G., Adefioye, O.A., Olowe, A.O., Opaleye, O.O., Bolaji, O.S., Ojurongbe. T.A. (2014): Soil Transmitted Helminths Infections among Primary School Children in Ile-Ife, South-West Nigeria: A cross-sectional study. *Int. Res. J. Med. Sci.* 2 (1): 6 – 10
 17. Otubanjo, O.A. and Mafe, M.A. (2002): Control of parasitic diseases of poverty: an overview of the Nigerian situation. *Nigerian Journal of Parasitology*, 1(1): 1-24.
 18. Steinmann P., Keizer J. and Bos R. (2006): Schistosomiasis and water resources development: Systematic Review, Meta-analysis, and estimates of people at risk. *The Lancet of Infectious Diseases*. 6:411-425
 19. Umoh, V.I, Okafor, C and Galadima, M. (2001): Contamination by helminths of vegetables cultivated on land irrigated with urban waste water in Zaria and Kaduna, Nigeria. *The Nigerian Journal of Parasitology*, 22(1&2): 95-104.
 20. WHO (2015) Investing to overcome the global impact of neglected tropical diseases; 3rd WHO report on Neglected Tropical Diseases.
 21. WHO (2011): Helminth Control in school-age children: a guide for managers of control programmes. 2nd edition. Geneva
 22. WHO (2010): Operational guide to mapping of Schistosomiasis and soil transmitted helminthiasis and evaluation of control programmes, WHO Geneva
 23. WHO (1991): *Basic Laboratory methods in medical parasitology* plates 1-9
 24. Worrell C, Davis S, Wiegand R, Lopez G, Odero K, (2013): Water, sanitation, and hygiene-related risk factors for soil-transmitted helminth infection in urban school- and pre-school-aged children in Kibera, Nairobi 2013; ASTMH 61st Annual Meeting; 11–15 November 2013, Atlanta.

APPENDIX

Appendix 1: Coordinated Mapping of Schistosomiasis & STH in Nigeria: Project Organogram



Appendix 2: Survey Teams

FMOH

Dr Bridget Okoeguale
Dr. Onyebuchi Uwaezuoke
Mrs I. Anagbogou
Dr. Sani Gwarzo
Dr. Evelyn Ngige
Dr. Jubril Suleiman
Dr. Obiageli J. Nebe
Mr. Adeleke M. Balogun
Mrs. Veronica S. Augustine
Dr. Nicolas Olobio
Dr. Uzoma Nwankwo
Mr. Nwoye Ikenna

Mr. Jacob Solomon M
Dr. Chukwudozie Chukwunonso
Mr. Adegunju Rotimi
Mr. Daguleng Ibrahim
Mrs. Rita O. Urude
Mr. Emmanuel Ibeku.C.
Miss. Omolola Iwayemi Olojede
Dr. Daramola Olakunle
Mrs. Nwafor Leticia
Mr. Shilo Paul
Mr. Afe Lawrence
Mr. Ren Whyte

NGDO TEAM

Dr. Sunday Isiyaku
Mr. Chris Ogoshi
Dr. (Mrs.) Francisca Olamiju
Dr (Mrs.) Ima Chima
Mr. Ben Nwobi
Mr. Adamani William
Mr. Christian Nwosu
Mr. Victor Osatogbe
Mrs. Jamila Abdel Adams
Mr. Jwanse Rinpan I.
Mrs. Esther Yusuf
Mr. Jacob Bako Emmanuel

Mr. Davou Dachomo
Mr. Nehemiah John
Mr. Aliyu Liman
Mr. Mba Ifeanyi Collins
Mr. Dennis Abrack
Mr. Obot. S. Samuel
Mr. Chidera Ukwunna
Mr. Stephen Abari
Mr. Sylvester Edet
Mr. Dinshiya Joel
Mr. Ages Atok

IN- COUNTRY CONSULTANTS

Prof. Dakul Anthony
Prof. Florence Nduka
Prof. Adenike. Abiose
Dr. Grace O. Adeoye
Dr. Njepuome Anthonia Ngozi

Dr. Amuga Gideon
Dr. Mwansat Georgina
Dr. Musa Sa'ad
Prof. O. Akogun
Dr. Francis Erah

JIGAWA STATE

Mr.A.U Abdulrahaman
Mr.Tijjani Muhammed
Mr.Aliyu UsmanTurali
Mr.Ismail Muhd. J
Mr.Halhiru Umar Imam
Mr.Bashir Mohammed I
Mr.Babangida Garba shitu
Mr.Kabiru Mohd
Mr.Abdulkadir Garba

Mr.Sani Hamisu
Mr.Rabiu Saidu
Mr.Abubakar Muhd.
Mr.Ibrahim Ado Ahmed
Mr.Basmah Sulaiman
Ms.Halisa Ashiru G.
Mr.Ibrahim Abdullahi
Mr.Aminu Yusuf Kazaure
Mr.Magaji Rabiu

KANO STATE

Mr. Danladi Tanko
Mr.Ibrahim Y Yahuza
Mr.Hamza Sule

Mr.Yunusa M. Aliyu
Mr. Abdullahi .T. Mohammed
Mrs. Zainab Isa Yalwa

Mrs. Amina Yusuf Ibrahim
Mr. Abdulkadir S. Wudilawa
Mr. Abdulwahab Umar
Mrs. Habibu Usman
Mr. Rabi'u Shehu Faiza
Mr. Saminu Ibrahim

KATSINA STATE

Mr. Samaila Mamman
Mr. Kabiru Dahiru
Mr. Yusuf Aliyu Runka
Mr. Ogbu Chinwe Mary
Mr. Bala Mohammed
Mr. Umaru Usman
Mr. Umaru Nasiru

LAGOS STATE

Mr. Bankole Munir A.
Mrs. Deborah Adepoju
Mr. Adeogun
Mr. Oshintayo
Mrs. Augusta
Mrs. Chinelo
Mr. Uche chidi
Mr. Olla
Mr. Olusanya

AKWA IBOM STATE

Dr. Veronica Itina
Mr. Basse E. Akpabio
Mrs. Blessing U. Akana – Ibiam
Mr. Akpan, Paul Nelson
Mr. Samuel Noah
Mr. Idorenyen N. U. Akpan
Mrs. Ruth A. Cletus

KOGI STATE

Mrs. Dorothy Amdife
Mr. Ojih A. Linus
Mr. Emmanuel Alaba
Mr. Abu Emmanuel
Mrs. Okenyi E. Stella
Mr. Usman Adulrazak
Mrs. Sheidu Sarah

KEBBI STATE

Mr. Attahiru Aliyu Aliero
Mr. Gimba Abrack
Mr. Musa Altine Algannara
Mr. Hassan Garba Jega
Mr. Aminu .A. Musa
Mr. Nafiu Farouk

Mr. Isa Auwal Shehu
Mr. Aminu Abubakar
Mr. Saifullahi Shehu
Mr. Hassan Ahmed
Mr. Danladi Adamu
Mr. Sirajo Sani

Mr. Ahmed Ibrahim
Mr. Kamilu Usman
Mr. Jinadu Murnai
Mr. Idris Abdilmajid
Mr. Bello Haliru
Mr. Bright Udeogu

Mrs. Duru Doris
Miss Elizabeth Ayodele
Mrs. Franca Opara
Mr. Adeseye
Mr. Aluko
Mr. Adejare
Mr. Okeke nneka
Mr. Akinwunmi

Mr. Esu Etim Okon
Mrs. Ekaette Tom Etuk
Mr. Ime O. Mr. Nwaha, Adiaha
Mr. John L. Umana
Mr. Nsebong John Dickson
Mr. Itina Gideon
Mrs. Christiana Okon

Mr. Yusuf M. B.
Mr. Okeme F. Jibril
Mr. Auwal Sani
Mrs. Adedoyin Y. M.
Mrs. Abiodun Florence
Mrs. Mark Rachel I.

Mr. Mohammed Muawiyah
Mrs. Kulu Umar
Mr. Aminu Samaila
Mrs. Halima Yakubu
Mr. Nasiru Mohammed Usman
Mr. Oluyemi Olusegun

Mr. Maksud Yusuf
Mr. Isa Abubakar Yahuri

Mr. Mohammadu Goru
Mr. Aliyu Bawa

STATE OF OSUN

Mr. Wahab Sule
Mr. Eso Jacob Olawole
Mr. Oladokun L. O
Mr. Braimah Rasheed
Mr. Oyelami Ebenezer
Mr. Oyrdokon Isiaka
Mr. Oyelere Bukola
Mr. Adewumni Olukemi

Mrs. Babalola E. F.
Mr. Adeleke Ayoade
Mr. Izu Micheal
Mr. Badejo Yinka
Mr. Sikiru
Mr. Oseni Kafaru
Mr. Oluwatosin C. O
Mr. J. T. Lagbaja

BAYELSA STATE

Obasi Johnson
Indakeme Oumo
Zifawei Kenneth
Sintei Timiondi B
Ebikebuna Alice
Kalial Idoumo

Ozu Eda
Igbelekumo Otamakiri
Andrew Grace
Omubo Grace
Bello Gregory
Chukwudebe Ifeoma

RIVERS STATE

Mr. Sammy. I. Eke
Mrs. Okwakpam Chika
Dr. Luke Anwuri
Mr. Ojirika Joy. C
Mr. Barisuka Nwibana
Mr. Kalio Future Greeson
Mrs. Oduku Heln
Mr. Nwayanwu Felix
Mr. Nkechi Ekeanyanma
Mr. Oghani Clementina
Mr. Labari Victoria N.

Mrs. Eleabu Gift. O
Mr. Ajih Jephthah
Mr. Mordecai Israel
Mr. Emmanuel. C. owhondah
Mr. Nwikina Komone
Mr. Ntor – ue Nnatah
Mr. Lysius. Z
Mr. Israel. R. Iorey
Mr. Eleabu Future
Mr. Monday Ikenga

OYO STATE

Mr. Alimi M. O
Mr. Taiwo M. O
Mr. Olaoti A. J
Mr. Adeniran Pius
Mr. Fagbemi Sunday
Mr. Adeniran Pius
Mr. Oluwasanmi O. A

Mr. Omotosho.A.A
Mr. Tolu Babalola
Mr. Sanusi Oluwasola
Mr. Rafiu Baasit
Mr. Ayodeji Jibril
Mr. Olaomi Olufunmito

YOBE STATE

Sani Zagam
Muhammed Mustapha
Sarki Mohammed Yusuf
Umar Audu Mohammed
Alh Maigari
Yusuf Mohammed
Haladu Isa Ahmed

Ayuba Goni Samson
Musa Galadima
Musa Madaki
Umar Muhmmad Baba
Fannami Gashi
Emmanuel Sarki Bulus
Mohammed Kachalla

Ahmed Alhaji Saidu

Toro Baba Lawan

FCT

Mrs Grace Ofodile
Mrs Josphine Igili
Mr. Nuhu Aliyu
Mr. Gideon Gwani
Mrs. Ugwu Ebere Vivial
Mr. Gideon Ogunmola
Mr. Innocent Attah
Mr. Sani Ibrahim

Mrs. Amina I. Michika
Mr. Idoma Peter. A
Mr. Ishaku Timlok Sule
Mr. Baba S. Dauda
Mr. Caleb k. Oga
Mr. Baba Aboki
Mr. Dauda
Mr. Abdullahi

BENUE STATE

Mrs. Betty Jande
Mr. Tarkigkir Henry. T
Mr. Onyilo. E. William
Mr. Ityoakoso Yaasa. J
Mr. Ityonzughul Joseph
Mr. Bernard Kumba
Mr. Najime Joseph

Mr. Idah David .I
Mrs. Margeret Iortim
Mrs. Kpeteh Esther. M
Mr. Moses Gbakaan
Mr. Abume Terzungwe
Mr. Thomas Terwase

NIGER STATE

Mr. Mallam Salisu
Mrs. Yusuf Aishatu
Mrs. Mariam Aminu
Mrs. Halima Ibrahim
Mr. Mohammed Takuma
Mr. Mohammed Bako
Mr. Musa Abubakar

Mr. Paul Elisha
Mr. Usman Ibrahim
Mrs. Ajah Doris
Mr. Taidi Gideon
Mr. .Mallam Salisu
Mr. Garuba Mohammed
Mr. Sylvanus Badien

BAUCHI STATE

Mr. Dahiru Mahmood
Mr. Abdullahi Abubakar
Mr. Nasiru Jibrin
Mr. Abdurahaman Danjuma
Mr. Yakubu Ahmed
Mr. Yunusa Adamu
Mr. Saleh Maimako
Mr. Rabiul Abdullahi
Mr. Sani Ibrahim
Mr. Mohammed Abubakar
Mr. Yakubu Chiroma

Mr. Sofian Audi
Mr. Aminu Hassan
Mr. Yusuf J. Sani
Mr. Damina Haruna
Mr. Bala Joshua Musa
Mr. Stephen Abari
Mrs. Franca Opara
Mr. Aboi Joseph Bodam
Mr. Coloumba
Ms. Ijeoma Achu
Mr. Manaseh Dakyen

TARABA STATE

Mr. Stephen S. Kaboson
Mr. Joel B. Dinshiya
Mr. Pajo Hikson
Mr. Mark Clement Adda
Mr. Daniel Mamman
Mr. Umar Musa Ibrahim

Mr. Nuhu Sambo
Mr. Nyako Finare
Mr. Salihu Bako
Mr. Alpha Dimas
Mr. Hammed Mogaji

EKITI STATE

Mrs. Alade .M .Irene
Mrs. Emmanuel. B .E
Mrs. Onuoha. O .F
Mrs. Tajudeen Wosilat
Ms. Adebias Bosede. B
Mrs. Fumilayo Da-Silva
Mrs. Oyebanji .O.I
Mr. Adesanmi Bayo
Mr. Akinwamide George
Mr. Ogundero G.O

Ms. Adeoumi Esther
Mr. Olofinlade Bode
Mrs, Ayeni Bukola
Mr. Akinyemi Taiwo
Mr. Aluko .K
Mr. Oluyole Oke
Mr. Obayemi. F
Mr. Oladipo Olusanya .S

KADUNA STATE

Mr. Alhassan Abdullahi
Mr. AbdulKarim Dauda
Mr. Adamu Mohammed
Mr. Haruna Damina
Mr. Mariam Abdulkadril
Mr. Jonah
Mr. Bature
Mr. Jonah M. Gado
Mr. Suleiman Ibrahim

Mr. Maikudi Yakubu
Mrs. Mary Victor
Mrs. Margret Kure
Mr. Mohammed
Mr. Shittu Ladan Abdullahi
Mr. Abba Mangai
Mr. Safiyanu Mohammed
Mr. Barau
Mr. Tanko

OTHERS

Mr. Famokun Adekunle
Mr. Ifeanyi Nwude
Mr. Pat-Nebe Okwudili. C.
Mr. Offor Solomon
Mr. Ajiji Joseph
Mr. Akpe Oluchi. E
Mrs. Blessing Dike
Mr. Eke Daniel Obinna

Mr. Dike Geoffrey. C
Mr. Saleh Usman
Miss. Obiora Chika
Mr. Emmanuel William
Mr. Obumneme Chigbo
Mr. Ezeobi. C. Daniel
Mrs. Funmilayo Olaniyan

Appendix 3: Data collection tools

CIFF School v3 (eng)

| Variable Name | Question Text | Saved Value | | | | | | | | | | | | |
|---------------|--|--|---|--------------------|---|------------------|---|----------------------|---|--------------------|---|------------------------------|---|---|
| meta | Hidden from user | | | | | | | | | | | | | |
| instanceID | Hidden from user | | | | | | | | | | | | | |
| SC0 | Date | User selected date | | | | | | | | | | | | |
| SC1 | Team Number | <table border="1"> <tr> <td>1</td> <td>Team 1</td> </tr> <tr> <td>2</td> <td>Team 2</td> </tr> <tr> <td>3</td> <td>Team 3</td> </tr> <tr> <td>4</td> <td>Team 4</td> </tr> <tr> <td>5</td> <td>Team 5</td> </tr> </table> | 1 | Team 1 | 2 | Team 2 | 3 | Team 3 | 4 | Team 4 | 5 | Team 5 | | |
| 1 | Team 1 | | | | | | | | | | | | | |
| 2 | Team 2 | | | | | | | | | | | | | |
| 3 | Team 3 | | | | | | | | | | | | | |
| 4 | Team 4 | | | | | | | | | | | | | |
| 5 | Team 5 | | | | | | | | | | | | | |
| SC4 | LGA Name | User entered text | | | | | | | | | | | | |
| SC5 | LGS Code | User entered text | | | | | | | | | | | | |
| SC6 | Community Name | User entered text | | | | | | | | | | | | |
| SC7 | School Name | User entered text | | | | | | | | | | | | |
| SC8 | Community/School Code | User entered text | | | | | | | | | | | | |
| SC8double | Community/School Code second entry | User entered text | | | | | | | | | | | | |
| SC9 | GPS | User captured location coordinates | | | | | | | | | | | | |
| SC12 | School Enrollment | User entered integer | | | | | | | | | | | | |
| SC13 | Number of Males Enrolled | User entered integer | | | | | | | | | | | | |
| SC14 | Number of Females Enrolled | User entered integer | | | | | | | | | | | | |
| SC16 | Number of Students Sampled | User entered integer | | | | | | | | | | | | |
| WS1 | Is there a source of drinking water in the school? | <table border="1"> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </table> | 0 | No | 1 | Yes | | | | | | | | |
| 0 | No | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | |
| WS2 | If yes, what type of water source? | <table border="1"> <tr> <td>1</td> <td>Unprotected spring</td> </tr> <tr> <td>2</td> <td>Protected spring</td> </tr> <tr> <td>3</td> <td>Unprotected dug well</td> </tr> <tr> <td>4</td> <td>Protected dug well</td> </tr> <tr> <td>5</td> <td>Hand pump/tube well/borehole</td> </tr> <tr> <td>6</td> <td>Surface water (river, dam, lake, stream, canal)</td> </tr> </table> | 1 | Unprotected spring | 2 | Protected spring | 3 | Unprotected dug well | 4 | Protected dug well | 5 | Hand pump/tube well/borehole | 6 | Surface water (river, dam, lake, stream, canal) |
| 1 | Unprotected spring | | | | | | | | | | | | | |
| 2 | Protected spring | | | | | | | | | | | | | |
| 3 | Unprotected dug well | | | | | | | | | | | | | |
| 4 | Protected dug well | | | | | | | | | | | | | |
| 5 | Hand pump/tube well/borehole | | | | | | | | | | | | | |
| 6 | Surface water (river, dam, lake, stream, canal) | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | | | | | | | |
|-----------|--|---|---|--------------------------------------|---|-----------------------|---|----------------------|----|----------------------|---|------------------------------|---|---|---|----------------------------------|---|----------------------|---|-------------------|----|----------------------|
| | | <table border="1"> <tbody> <tr> <td>7</td> <td>Public piped water/tap/standpipe</td> </tr> <tr> <td>8</td> <td>Rainwater collection</td> </tr> <tr> <td>9</td> <td>Sachet/Pure water</td> </tr> <tr> <td>10</td> <td>Other (specify next)</td> </tr> </tbody> </table> | 7 | Public piped water/tap/standpipe | 8 | Rainwater collection | 9 | Sachet/Pure water | 10 | Other (specify next) | | | | | | | | | | | | |
| 7 | Public piped water/tap/standpipe | | | | | | | | | | | | | | | | | | | | | |
| 8 | Rainwater collection | | | | | | | | | | | | | | | | | | | | | |
| 9 | Sachet/Pure water | | | | | | | | | | | | | | | | | | | | | |
| 10 | Other (specify next) | | | | | | | | | | | | | | | | | | | | | |
| WS2specif | Specify, other source of water | User entered text | | | | | | | | | | | | | | | | | | | | |
| WS3 | Are there sources of drinking water close to the school? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| WS4 | If yes, what type of water source? | <table border="1"> <tbody> <tr> <td>1</td> <td>Unprotected spring</td> </tr> <tr> <td>2</td> <td>Protected spring</td> </tr> <tr> <td>3</td> <td>Unprotected dug well</td> </tr> <tr> <td>4</td> <td>Protected dug well</td> </tr> <tr> <td>5</td> <td>Hand pump/tube well/borehole</td> </tr> <tr> <td>6</td> <td>Surface water (river, dam, lake, stream, canal)</td> </tr> <tr> <td>7</td> <td>Public piped water/tap/standpipe</td> </tr> <tr> <td>8</td> <td>Rainwater collection</td> </tr> <tr> <td>9</td> <td>Sachet/Pure water</td> </tr> <tr> <td>10</td> <td>Other (specify next)</td> </tr> </tbody> </table> | 1 | Unprotected spring | 2 | Protected spring | 3 | Unprotected dug well | 4 | Protected dug well | 5 | Hand pump/tube well/borehole | 6 | Surface water (river, dam, lake, stream, canal) | 7 | Public piped water/tap/standpipe | 8 | Rainwater collection | 9 | Sachet/Pure water | 10 | Other (specify next) |
| 1 | Unprotected spring | | | | | | | | | | | | | | | | | | | | | |
| 2 | Protected spring | | | | | | | | | | | | | | | | | | | | | |
| 3 | Unprotected dug well | | | | | | | | | | | | | | | | | | | | | |
| 4 | Protected dug well | | | | | | | | | | | | | | | | | | | | | |
| 5 | Hand pump/tube well/borehole | | | | | | | | | | | | | | | | | | | | | |
| 6 | Surface water (river, dam, lake, stream, canal) | | | | | | | | | | | | | | | | | | | | | |
| 7 | Public piped water/tap/standpipe | | | | | | | | | | | | | | | | | | | | | |
| 8 | Rainwater collection | | | | | | | | | | | | | | | | | | | | | |
| 9 | Sachet/Pure water | | | | | | | | | | | | | | | | | | | | | |
| 10 | Other (specify next) | | | | | | | | | | | | | | | | | | | | | |
| WS4specif | Specify, other source of drinking water close to school | User entered text | | | | | | | | | | | | | | | | | | | | |
| WS5 | Are there accessible water bodies close to the school? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| PL1 | Is there a latrine in the school? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| PL2 | Evidence of latrine usage observed? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| PL3 | What type of latrine is present? | <table border="1"> <tbody> <tr> <td>1</td> <td>Pit latrine without slab or open pit</td> </tr> <tr> <td>2</td> <td>Pit latrine with slab</td> </tr> </tbody> </table> | 1 | Pit latrine without slab or open pit | 2 | Pit latrine with slab | | | | | | | | | | | | | | | | |
| 1 | Pit latrine without slab or open pit | | | | | | | | | | | | | | | | | | | | | |
| 2 | Pit latrine with slab | | | | | | | | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | | | | | | | |
|--------------------|---|--|---|--------------------------------------|---|-----------------------|---|----------------------|----|------------------------------|---|------------------------------|---|---|---|----------------------------------|---|----------------------|---|-------------------|----|------------------------------|
| | | <table border="1"> <tbody> <tr> <td>7</td> <td>Public piped water/tap/standpipe</td> </tr> <tr> <td>8</td> <td>Rainwater collection</td> </tr> <tr> <td>9</td> <td>Sachet/Pure water</td> </tr> <tr> <td>10</td> <td>Other (<i>specify</i> next)</td> </tr> </tbody> </table> | 7 | Public piped water/tap/standpipe | 8 | Rainwater collection | 9 | Sachet/Pure water | 10 | Other (<i>specify</i> next) | | | | | | | | | | | | |
| 7 | Public piped water/tap/standpipe | | | | | | | | | | | | | | | | | | | | | |
| 8 | Rainwater collection | | | | | | | | | | | | | | | | | | | | | |
| 9 | Sachet/Pure water | | | | | | | | | | | | | | | | | | | | | |
| 10 | Other (<i>specify</i> next) | | | | | | | | | | | | | | | | | | | | | |
| WS2 <i>specify</i> | <i>Specify</i> , other source of water | User entered text | | | | | | | | | | | | | | | | | | | | |
| WS3 | Are there sources of drinking water close to the school? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| WS4 | If <i>yes</i> , what <i>type</i> of water source? | <table border="1"> <tbody> <tr> <td>1</td> <td>Unprotected spring</td> </tr> <tr> <td>2</td> <td>Protected spring</td> </tr> <tr> <td>3</td> <td>Unprotected dug well</td> </tr> <tr> <td>4</td> <td>Protected dug well</td> </tr> <tr> <td>6</td> <td>Hand pump/tube well/borehole</td> </tr> <tr> <td>6</td> <td>Surface water (river, dam, lake, stream, canal)</td> </tr> <tr> <td>7</td> <td>Public piped water/tap/standpipe</td> </tr> <tr> <td>8</td> <td>Rainwater collection</td> </tr> <tr> <td>9</td> <td>Sachet/Pure water</td> </tr> <tr> <td>10</td> <td>Other (<i>specify</i> next)</td> </tr> </tbody> </table> | 1 | Unprotected spring | 2 | Protected spring | 3 | Unprotected dug well | 4 | Protected dug well | 6 | Hand pump/tube well/borehole | 6 | Surface water (river, dam, lake, stream, canal) | 7 | Public piped water/tap/standpipe | 8 | Rainwater collection | 9 | Sachet/Pure water | 10 | Other (<i>specify</i> next) |
| 1 | Unprotected spring | | | | | | | | | | | | | | | | | | | | | |
| 2 | Protected spring | | | | | | | | | | | | | | | | | | | | | |
| 3 | Unprotected dug well | | | | | | | | | | | | | | | | | | | | | |
| 4 | Protected dug well | | | | | | | | | | | | | | | | | | | | | |
| 6 | Hand pump/tube well/borehole | | | | | | | | | | | | | | | | | | | | | |
| 6 | Surface water (river, dam, lake, stream, canal) | | | | | | | | | | | | | | | | | | | | | |
| 7 | Public piped water/tap/standpipe | | | | | | | | | | | | | | | | | | | | | |
| 8 | Rainwater collection | | | | | | | | | | | | | | | | | | | | | |
| 9 | Sachet/Pure water | | | | | | | | | | | | | | | | | | | | | |
| 10 | Other (<i>specify</i> next) | | | | | | | | | | | | | | | | | | | | | |
| WS4 <i>specify</i> | <i>Specify</i> , other source of drinking water close to school | User entered text | | | | | | | | | | | | | | | | | | | | |
| WS5 | Are there accessible water bodies close to the school? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| PL1 | Is there a latrine in the school? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| PL2 | Evidence of latrine usage observed? | <table border="1"> <tbody> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </tbody> </table> | 0 | No | 1 | Yes | | | | | | | | | | | | | | | | |
| 0 | No | | | | | | | | | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | | | | | | | | | |
| PL3 | What <i>type</i> of latrine is present? | <table border="1"> <tbody> <tr> <td>1</td> <td>Pit latrine without slab or open pit</td> </tr> <tr> <td>2</td> <td>Pit latrine with slab</td> </tr> </tbody> </table> | 1 | Pit latrine without slab or open pit | 2 | Pit latrine with slab | | | | | | | | | | | | | | | | |
| 1 | Pit latrine without slab or open pit | | | | | | | | | | | | | | | | | | | | | |
| 2 | Pit latrine with slab | | | | | | | | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | |
|-------------|--|---|---|--|---|--|----|---|---|--|---|---|---|----------------|
| | | <table border="1"> <tr> <td>3</td> <td>Ventilated improved pit latrine (VIP)</td> </tr> <tr> <td>4</td> <td>Flush or pour flush toilet</td> </tr> <tr> <td>99</td> <td>Other</td> </tr> </table> | 3 | Ventilated improved pit latrine (VIP) | 4 | Flush or pour flush toilet | 99 | Other | | | | | | |
| 3 | Ventilated improved pit latrine (VIP) | | | | | | | | | | | | | |
| 4 | Flush or pour flush toilet | | | | | | | | | | | | | |
| 99 | Other | | | | | | | | | | | | | |
| PL4 | What is the condition of the latrine? | <table border="1"> <tr> <td>0</td> <td>Poor (Presence of flies, offensive odour and visible stool on floor, absence of roof/door)</td> </tr> <tr> <td>1</td> <td>Fair (Presence of roof/door but dirty floor)</td> </tr> <tr> <td>3</td> <td>Moderate (clean, absence of roof or door)</td> </tr> <tr> <td>4</td> <td>Good (Clean, odourless, no flies, presence of roof and door)</td> </tr> <tr> <td>6</td> <td>Excellent (Very clean, odourless, presence of roof, door and availability of water)</td> </tr> </table> | 0 | Poor (Presence of flies, offensive odour and visible stool on floor, absence of roof/door) | 1 | Fair (Presence of roof/door but dirty floor) | 3 | Moderate (clean, absence of roof or door) | 4 | Good (Clean, odourless, no flies, presence of roof and door) | 6 | Excellent (Very clean, odourless, presence of roof, door and availability of water) | | |
| 0 | Poor (Presence of flies, offensive odour and visible stool on floor, absence of roof/door) | | | | | | | | | | | | | |
| 1 | Fair (Presence of roof/door but dirty floor) | | | | | | | | | | | | | |
| 3 | Moderate (clean, absence of roof or door) | | | | | | | | | | | | | |
| 4 | Good (Clean, odourless, no flies, presence of roof and door) | | | | | | | | | | | | | |
| 6 | Excellent (Very clean, odourless, presence of roof, door and availability of water) | | | | | | | | | | | | | |
| PL5 | Is there water or Tissue for use after defecating? | <table border="1"> <tr> <td>1</td> <td>Always</td> </tr> <tr> <td>2</td> <td>Sometimes</td> </tr> <tr> <td>3</td> <td>Never</td> </tr> <tr> <td>9</td> <td>Other, Specify</td> </tr> </table> | 1 | Always | 2 | Sometimes | 3 | Never | 9 | Other, Specify | | | | |
| 1 | Always | | | | | | | | | | | | | |
| 2 | Sometimes | | | | | | | | | | | | | |
| 3 | Never | | | | | | | | | | | | | |
| 9 | Other, Specify | | | | | | | | | | | | | |
| PL5specify | Specify, other. Is there water or Tissue for use after defecating? | User entered text | | | | | | | | | | | | |
| PL6 | Is there provision for hand washing after toilet use? | <table border="1"> <tr> <td>0</td> <td>No</td> </tr> <tr> <td>1</td> <td>Yes</td> </tr> </table> | 0 | No | 1 | Yes | | | | | | | | |
| 0 | No | | | | | | | | | | | | | |
| 1 | Yes | | | | | | | | | | | | | |
| PL7 | What type of hand washing facilities? | <table border="1"> <tr> <td>0</td> <td>No water</td> </tr> <tr> <td>1</td> <td>Water only</td> </tr> <tr> <td>2</td> <td>Water and soap</td> </tr> <tr> <td>3</td> <td>Water, soap, and non-disposable napkin</td> </tr> <tr> <td>4</td> <td>Water, soap, and disposable napkin</td> </tr> <tr> <td>9</td> <td>Other, specify</td> </tr> </table> | 0 | No water | 1 | Water only | 2 | Water and soap | 3 | Water, soap, and non-disposable napkin | 4 | Water, soap, and disposable napkin | 9 | Other, specify |
| 0 | No water | | | | | | | | | | | | | |
| 1 | Water only | | | | | | | | | | | | | |
| 2 | Water and soap | | | | | | | | | | | | | |
| 3 | Water, soap, and non-disposable napkin | | | | | | | | | | | | | |
| 4 | Water, soap, and disposable napkin | | | | | | | | | | | | | |
| 9 | Other, specify | | | | | | | | | | | | | |
| PL7specify | Specify, other. What type of hand washing facilities? | User entered text | | | | | | | | | | | | |
| SchoolNotes | Additional notes (optional) | User entered text | | | | | | | | | | | | |

CIFF Student v4 (eng)

| Variable Name | Question Text | Saved Value | | | | | | | | | | | | |
|---------------|--|---|---|----------------------|---|------------------------|---|----------------------------|---|----------------|---|-----------------|----|------------|
| meta | Hidden from user | | | | | | | | | | | | | |
| instanceID | Hidden from user | | | | | | | | | | | | | |
| ST0 | Date | User selected date | | | | | | | | | | | | |
| ST1 | Team Number | <table border="1"> <tr> <td>1</td> <td>Team 1</td> </tr> <tr> <td>2</td> <td>Team 2</td> </tr> <tr> <td>3</td> <td>Team 3</td> </tr> <tr> <td>4</td> <td>Team 4</td> </tr> <tr> <td>5</td> <td>Team 5</td> </tr> </table> | 1 | Team 1 | 2 | Team 2 | 3 | Team 3 | 4 | Team 4 | 5 | Team 5 | | |
| 1 | Team 1 | | | | | | | | | | | | | |
| 2 | Team 2 | | | | | | | | | | | | | |
| 3 | Team 3 | | | | | | | | | | | | | |
| 4 | Team 4 | | | | | | | | | | | | | |
| 5 | Team 5 | | | | | | | | | | | | | |
| ST6 | Student UID | User entered text | | | | | | | | | | | | |
| ST6double | Student UID second entry | User entered text | | | | | | | | | | | | |
| ST6 | Name | User entered text | | | | | | | | | | | | |
| ST7 | Age | User entered integer | | | | | | | | | | | | |
| ST8 | Sex | <table border="1"> <tr> <td>F</td> <td>Female</td> </tr> <tr> <td>M</td> <td>Male</td> </tr> </table> | F | Female | M | Male | | | | | | | | |
| F | Female | | | | | | | | | | | | | |
| M | Male | | | | | | | | | | | | | |
| ST9 | When <i>you</i> are at school where do <i>you</i> usually go to URINATE? | <table border="1"> <tr> <td>0</td> <td>In the school toilet</td> </tr> <tr> <td>1</td> <td>Around school compound</td> </tr> <tr> <td>2</td> <td>Outside of school compound</td> </tr> <tr> <td>3</td> <td>I wait/hold it</td> </tr> <tr> <td>9</td> <td>Others, specify</td> </tr> <tr> <td>99</td> <td>Don't know</td> </tr> </table> | 0 | In the school toilet | 1 | Around school compound | 2 | Outside of school compound | 3 | I wait/hold it | 9 | Others, specify | 99 | Don't know |
| 0 | In the school toilet | | | | | | | | | | | | | |
| 1 | Around school compound | | | | | | | | | | | | | |
| 2 | Outside of school compound | | | | | | | | | | | | | |
| 3 | I wait/hold it | | | | | | | | | | | | | |
| 9 | Others, specify | | | | | | | | | | | | | |
| 99 | Don't know | | | | | | | | | | | | | |
| ST9specify | Specify, other. When <i>you</i> are at school where do <i>you</i> usually go to URINATE? | User entered text | | | | | | | | | | | | |
| ST10 | When <i>you</i> are at school where do <i>you</i> usually go to DEFECATE? | <table border="1"> <tr> <td>0</td> <td>In the school toilet</td> </tr> <tr> <td>1</td> <td>Around school compound</td> </tr> <tr> <td>2</td> <td>Outside of school compound</td> </tr> <tr> <td>3</td> <td>I wait/hold it</td> </tr> <tr> <td>9</td> <td>Others, specify</td> </tr> </table> | 0 | In the school toilet | 1 | Around school compound | 2 | Outside of school compound | 3 | I wait/hold it | 9 | Others, specify | | |
| 0 | In the school toilet | | | | | | | | | | | | | |
| 1 | Around school compound | | | | | | | | | | | | | |
| 2 | Outside of school compound | | | | | | | | | | | | | |
| 3 | I wait/hold it | | | | | | | | | | | | | |
| 9 | Others, specify | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | | | |
|------------|--|--|---|---------|---|---------|---|---------|---|----------------|---|----------------|---|---------|---|----------|---|-------|
| | | 99 Don't know | | | | | | | | | | | | | | | | |
| ST10specif | Specify, other. When you are at school where do you usually go to DEFECATE? | User entered text | | | | | | | | | | | | | | | | |
| ST11 | If there are water bodies, which of the following activities do you perform? | <table border="1"> <tr><td>1</td><td>Bathing</td></tr> <tr><td>2</td><td>Washing</td></tr> <tr><td>3</td><td>Fishing</td></tr> <tr><td>4</td><td>Crossing water</td></tr> <tr><td>5</td><td>Fetching water</td></tr> <tr><td>6</td><td>Playing</td></tr> <tr><td>7</td><td>Swimming</td></tr> <tr><td>9</td><td>Other</td></tr> </table> | 1 | Bathing | 2 | Washing | 3 | Fishing | 4 | Crossing water | 5 | Fetching water | 6 | Playing | 7 | Swimming | 9 | Other |
| 1 | Bathing | | | | | | | | | | | | | | | | | |
| 2 | Washing | | | | | | | | | | | | | | | | | |
| 3 | Fishing | | | | | | | | | | | | | | | | | |
| 4 | Crossing water | | | | | | | | | | | | | | | | | |
| 5 | Fetching water | | | | | | | | | | | | | | | | | |
| 6 | Playing | | | | | | | | | | | | | | | | | |
| 7 | Swimming | | | | | | | | | | | | | | | | | |
| 9 | Other | | | | | | | | | | | | | | | | | |
| ST11specif | Specify, other. If there are water bodies, do you perform any of the following activities? | User entered text | | | | | | | | | | | | | | | | |
| ST12 | Additional notes (optional) | User entered text | | | | | | | | | | | | | | | | |

CIFF Stool sample v4 (eng)

| Variable Name | Question Text | Saved Value | | | | | | | | | | |
|---------------|--|--|---|--------|---|--------|---|--------|---|--------|---|--------|
| meta | Hidden from user | | | | | | | | | | | |
| instanceID | Hidden from user | | | | | | | | | | | |
| STL0 | Date | User selected date | | | | | | | | | | |
| STL1 | Team Number | <table border="1"> <tbody> <tr> <td>1</td> <td>Team 1</td> </tr> <tr> <td>2</td> <td>Team 2</td> </tr> <tr> <td>3</td> <td>Team 3</td> </tr> <tr> <td>4</td> <td>Team 4</td> </tr> <tr> <td>6</td> <td>Team 6</td> </tr> </tbody> </table> | 1 | Team 1 | 2 | Team 2 | 3 | Team 3 | 4 | Team 4 | 6 | Team 6 |
| 1 | Team 1 | | | | | | | | | | | |
| 2 | Team 2 | | | | | | | | | | | |
| 3 | Team 3 | | | | | | | | | | | |
| 4 | Team 4 | | | | | | | | | | | |
| 6 | Team 6 | | | | | | | | | | | |
| STL5 | Student UID | User entered text | | | | | | | | | | |
| STL5double | Student UID second entry | User entered text | | | | | | | | | | |
| STL6 | <i>Ascaris lumbricoides</i> eggs/slide | User entered integer | | | | | | | | | | |
| STL7 | <i>Trichuris trichiura</i> eggs/slide | User entered integer | | | | | | | | | | |
| STL8 | Hookworm eggs/slide | User entered integer | | | | | | | | | | |
| STL9 | <i>Schistosoma mansoni</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST10 | <i>T. Orientalis</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST11 | <i>E. histolytica</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST12 | <i>E. Vermicularis</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST13 | <i>D.latum</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST14 | <i>T.spp</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST16 | <i>F. Hepatica</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST16 | <i>Hymenolepis nana</i> eggs/slide | User entered integer | | | | | | | | | | |
| ST17 | <i>E.coli</i> | User entered integer | | | | | | | | | | |
| ST18 | Additional notes (optional) | User entered text | | | | | | | | | | |


CIFF Urine sample v3 (eng)

| Variable Name | Question Text | Saved Value | | | | | | | | | | |
|---------------|-----------------------------|--|---|--------|---|--------------|---|-------------|---|-------------------|---|--------------------|
| meta | Hidden from user | | | | | | | | | | | |
| instanceID | Hidden from user | | | | | | | | | | | |
| UR0 | Date | User selected date | | | | | | | | | | |
| UR1 | Team Number | <table border="1"> <tbody> <tr> <td>1</td> <td>Team 1</td> </tr> <tr> <td>2</td> <td>Team 2</td> </tr> <tr> <td>3</td> <td>Team 3</td> </tr> <tr> <td>4</td> <td>Team 4</td> </tr> <tr> <td>5</td> <td>Team 5</td> </tr> </tbody> </table> | 1 | Team 1 | 2 | Team 2 | 3 | Team 3 | 4 | Team 4 | 5 | Team 5 |
| 1 | Team 1 | | | | | | | | | | | |
| 2 | Team 2 | | | | | | | | | | | |
| 3 | Team 3 | | | | | | | | | | | |
| 4 | Team 4 | | | | | | | | | | | |
| 5 | Team 5 | | | | | | | | | | | |
| UR5 | Student UID | User entered text | | | | | | | | | | |
| UR5double | Student UID second entry | User entered text | | | | | | | | | | |
| UR6 | Urine color | <table border="1"> <tbody> <tr> <td>1</td> <td>Bloody</td> </tr> <tr> <td>2</td> <td>Amber/Cloudy</td> </tr> <tr> <td>3</td> <td>Amber/Clear</td> </tr> <tr> <td>4</td> <td>Pale yellow/Clear</td> </tr> <tr> <td>5</td> <td>Pale yellow/Cloudy</td> </tr> </tbody> </table> | 1 | Bloody | 2 | Amber/Cloudy | 3 | Amber/Clear | 4 | Pale yellow/Clear | 5 | Pale yellow/Cloudy |
| 1 | Bloody | | | | | | | | | | | |
| 2 | Amber/Cloudy | | | | | | | | | | | |
| 3 | Amber/Clear | | | | | | | | | | | |
| 4 | Pale yellow/Clear | | | | | | | | | | | |
| 5 | Pale yellow/Cloudy | | | | | | | | | | | |
| UR7 | HEM Urine Dipstick | <table border="1"> <tbody> <tr> <td>0</td> <td>Neg</td> </tr> <tr> <td>1</td> <td>Trace</td> </tr> <tr> <td>2</td> <td>+</td> </tr> <tr> <td>3</td> <td>++</td> </tr> <tr> <td>4</td> <td>+++</td> </tr> </tbody> </table> | 0 | Neg | 1 | Trace | 2 | + | 3 | ++ | 4 | +++ |
| 0 | Neg | | | | | | | | | | | |
| 1 | Trace | | | | | | | | | | | |
| 2 | + | | | | | | | | | | | |
| 3 | ++ | | | | | | | | | | | |
| 4 | +++ | | | | | | | | | | | |
| UR8 | No of eggs per 10mls | User entered integer | | | | | | | | | | |
| UR9 | Additional notes (optional) | User entered text | | | | | | | | | | |

CIFF error report v2 (eng)

| Variable Name | Question Text | Saved Value | | | | | | | | |
|---------------|---|--|---------|---------|--------|--------|-------|-------|-------|-------|
| meta | Hidden from user | | | | | | | | | |
| instanceID | Hidden from user | | | | | | | | | |
| Error0 | Date | User selected date | | | | | | | | |
| Error1 | Team Number | User entered integer | | | | | | | | |
| Error2 | Choose which survey has the error | <table border="1"> <tbody> <tr> <td>Student</td> <td>Student</td> </tr> <tr> <td>School</td> <td>School</td> </tr> <tr> <td>Urine</td> <td>Urine</td> </tr> <tr> <td>Stool</td> <td>Stool</td> </tr> </tbody> </table> | Student | Student | School | School | Urine | Urine | Stool | Stool |
| Student | Student | | | | | | | | | |
| School | School | | | | | | | | | |
| Urine | Urine | | | | | | | | | |
| Stool | Stool | | | | | | | | | |
| Error3 | Details for the error you are reporting | User entered text | | | | | | | | |

Appendix 4: Flow Diagram of the Android Based Electronic Data Management



Appendix 5: Ethical Clearance



National Health Research Ethics Committee of Nigeria (NHREC)

Promoting Highest Ethical and Scientific Standards
for Health Research in Nigeria



Federal Ministry of Health

NHREC Protocol Number NHREC/01/01/2007-18/07/2014
NHREC Approval Number NHREC/01/01/2007-25/11/2014c
Date: 25th November, 2014

RE: INTEGRATED EPIDEMIOLOGICAL MAPPING AND BASELINE SURVEY OF NTDS (LYMPHATIC FILARIASIS, SCHISTOSOMIASIS AND SOIL TRANSMITTED HELMINTHS) IN NIGERIA

Health Research Ethics Committee (HREC) assigned number: NHREC/01/01/2007

Name of Principal Investigator: Dr. Uwaezuoke Onyebuchi

Address of Investigator: Department of Public Health
Federal Ministry of Health
Cell Phone: +2348089023030
E-mail: happywaez@yahoo.com

Date of receipt of valid application: 18-07-2014

Date when final determination of research was made: 25-11-2014

Notice of Expedited Review and Approval

This is to inform you that the research described in the submitted protocol, the consent forms, other participant information materials have been reviewed and given expedited committee approval by the National Health Research Ethics Committee.

This approval dates remain as in the initial approval from 25/11/2014 to 24/11/2015. If there is delay in starting the research, please inform the HREC so that the dates of approval can be adjusted accordingly. Note that no participant accrual or activity related to this research may be conducted outside of these dates. *All informed consent forms used in this study must carry the HREC assigned number and duration of HREC approval of the study.* If this is a multi-year research, endeavour to submit your annual report to the HREC early in order to obtain renewal of your approval and avoid disruption of your research.

The National Code for Health Research Ethics requires you to comply with all institutional guidelines, rules and regulations and with the tenets of the Code including ensuring that all adverse events are reported promptly to the HREC. No changes are permitted in the research without prior approval by the HREC except in circumstances outlined in the Code. The HREC reserves the right to conduct compliance visit your research site without previous notification.

Signed

Clement Adebamowo BMChB Hons (Jos), FWACS, FACS, DSc (Harvard)
Chairman, National Health Research Ethics Committee of Nigeria (NHREC)

Department of Health Planning, Research & Statistics
Federal Ministry of Health
11th Floor, Federal Secretariat Complex Phase III
Ahmadu Bello Way, Abuja

Tel: +234-09-523-8367
E-mail: chairman@nhrec.net, secretary@nhrec.net,
deskofficer@nhrec.net,
URL: <http://www.nhrec.net>

Appendix 6: Treatment guideline for schistosomiasis/STH

| Category | Disease | Prevalence (by parasitological methods) | Action | |
|---------------|-----------------|--|--|--|
| High-risk | Schistosomiasis | ≥50% | Treat all school-age children once per year | Also treat adults considered to be at risk |
| Moderate-risk | Schistosomiasis | ≥10% but <50% | Treat all school-age children once every two years | Also treat adults considered to be at risk |
| Low-risk | Schistosomiasis | <10% | Treat all school-age children twice during their primary schooling | Praziquantel should be available in dispensaries and clinics |
| High-risk | STH | ≥50% | Treat all school-age children twice per year | Also treat: <ul style="list-style-type: none"> • preschool children • women of childbearing age, including pregnant women in the 2nd and 3rd trimesters and lactating women • adults in high risk occupations |
| Low risk | STH | ≥20% but <50% | Treat all school-age children once per year | Also treat: <ul style="list-style-type: none"> • preschool children • women of childbearing age, including pregnant women in the 2nd and 3rd trimesters and lactating women • adults in high risk occupations |
| | STH | <20% | Case-by case management of those infected | |

Source: WHO/CTD/SIP/98.1

Appendix 7: LGAs Requiring Interventions

| State | LGA Name | Schistosomiasis | | | STH | | |
|-----------|-------------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Akwa-Ibom | Itu | - | - | - | - | ✓ | - |
| Akwa-Ibom | Uruan | - | - | - | - | ✓ | - |
| Akwa-Ibom | Urue-Offong/Oruko | - | - | - | - | ✓ | - |
| Akwa-Ibom | Ibesikpo Asutan | - | - | - | - | ✓ | - |
| Akwa-Ibom | Ini | - | - | - | - | ✓ | - |
| Akwa-Ibom | Nsit-Ibom | - | - | - | ✓ | - | - |
| Akwa-Ibom | Uyo | - | - | - | ✓ | - | - |
| Akwa-Ibom | Oruk Anam | - | - | - | - | ✓ | - |
| Akwa-Ibom | Ikot Ekpene | - | - | - | ✓ | - | - |
| Akwa-Ibom | Ikono | - | - | - | ✓ | - | - |
| Akwa-Ibom | Essien Udim | - | - | - | ✓ | - | - |
| Akwa-Ibom | Ika | - | - | - | ✓ | - | - |
| Akwa-Ibom | Abak | - | - | - | - | ✓ | - |
| Akwa-Ibom | Ibiono Ibom | - | - | - | - | ✓ | - |
| Akwa-Ibom | Etim Ekpo | - | - | - | ✓ | - | - |
| Akwa-Ibom | Obot Akara | - | - | - | - | ✓ | - |
| Akwa-Ibom | Udung-Uko | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Mbo | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Ibena | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Okobo | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Oron | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Ikot Abasi | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Onna | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Eastern Obolo | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Mkpat-Enin | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Esit Eket | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Ukanafun | ✓ | - | - | ✓ | - | - |
| Akwa-Ibom | Nsit-Atai | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Nsit-Ubium | ✓ | - | - | - | ✓ | - |
| Akwa-Ibom | Etinan | ✓ | - | - | ✓ | - | - |
| Akwa-Ibom | Eket | ✓ | - | - | - | ✓ | - |
| Bauchi | Warji | - | ✓ | - | - | - | ✓ |
| Bauchi | Bauchi | - | ✓ | - | - | - | ✓ |
| Bauchi | Shira | - | ✓ | - | - | - | ✓ |
| Bauchi | Misau | - | - | - | - | - | ✓ |
| Bauchi | Ningi | - | ✓ | - | - | - | ✓ |
| Bauchi | Alkaleri | - | ✓ | - | - | - | ✓ |
| Bauchi | Itas_Gadua | - | ✓ | - | - | - | ✓ |
| Bauchi | Gamawa | - | ✓ | - | - | - | ✓ |
| Bauchi | Tafawa Balewa | - | ✓ | - | - | - | ✓ |
| Bauchi | Zaki | - | ✓ | - | - | - | ✓ |

| State | LGA Name | Schistosomiasis | | | STH | | |
|-------------|-------------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Bauchi | Bogoro | - | ✓ | - | - | - | ✓ |
| Bauchi | Ganjuwa | - | ✓ | - | - | - | ✓ |
| Bauchi | Katagum | ✓ | - | - | - | - | ✓ |
| Bauchi | Kirfi | ✓ | - | - | - | - | ✓ |
| Bauchi | Giade | ✓ | - | - | - | - | ✓ |
| Bauchi | Jama'are | ✓ | - | - | - | - | ✓ |
| Bauchi | Darazo | ✓ | - | - | - | - | ✓ |
| Bauchi | Dambam | ✓ | - | - | - | - | ✓ |
| Bauchi | Dass | ✓ | - | - | - | - | ✓ |
| Bauchi | Toro | ✓ | - | - | - | - | ✓ |
| Bayelsa | Brass | - | - | - | - | ✓ | - |
| Bayelsa | Nembe | - | - | - | - | - | ✓ |
| Bayelsa | Southern Ijaw | ✓ | - | - | - | ✓ | - |
| Bayelsa | Ekeremor | ✓ | - | - | ✓ | - | - |
| Bayelsa | Yenagoa | ✓ | - | - | ✓ | - | - |
| Bayelsa | Ogbia | ✓ | - | - | ✓ | - | - |
| Bayelsa | Sagbama | ✓ | - | - | - | - | ✓ |
| Bayelsa | Kolokuma/Opokuma | ✓ | - | - | - | - | ✓ |
| Benue | Gwer East | - | ✓ | - | ✓ | - | - |
| Benue | Agatu | ✓ | - | - | ✓ | - | - |
| Benue | Guma | - | ✓ | - | ✓ | - | - |
| Benue | Oju | - | ✓ | - | ✓ | - | - |
| Benue | Gwer West | - | ✓ | - | ✓ | - | - |
| Benue | Ukum | - | ✓ | - | ✓ | - | - |
| Benue | Apa | - | ✓ | - | ✓ | - | - |
| Benue | Ado | ✓ | - | - | ✓ | - | - |
| Benue | Vandeikya | - | ✓ | - | ✓ | - | - |
| Benue | Obi | - | ✓ | - | ✓ | - | - |
| Benue | Konshisha | ✓ | - | - | ✓ | - | - |
| Benue | Ogbadibo | ✓ | - | - | ✓ | - | - |
| Benue | Ohimini | ✓ | - | - | ✓ | - | - |
| Benue | Tarka | - | ✓ | - | ✓ | - | - |
| Cross River | Bakasi | - | - | - | ✓ | - | - |
| Cross River | Akpabuyo | - | - | - | ✓ | - | - |
| Cross River | Biase | - | - | - | ✓ | - | - |
| Cross River | Akamkpa | - | - | - | ✓ | - | - |
| Cross River | Calabar South | - | - | - | ✓ | - | - |
| Cross River | Calabar Municipal | - | - | - | ✓ | - | - |
| Cross River | Obanliku | - | ✓ | - | - | - | ✓ |
| Cross River | Ogoja | - | ✓ | - | - | - | ✓ |
| Cross River | Obudu | - | ✓ | - | - | - | ✓ |
| Cross River | Yala | - | ✓ | - | - | - | ✓ |
| Cross River | Yakurr | ✓ | - | - | ✓ | - | - |

| State | LGA Name | Schistosomiasis | | | STH | | |
|-------------|------------------------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Cross River | Odukpani | ✓ | - | - | - | ✓ | - |
| Cross River | Etung | ✓ | - | - | ✓ | - | - |
| Cross River | Ikom | ✓ | - | - | - | - | ✓ |
| Cross River | Boki | ✓ | - | - | - | - | ✓ |
| Cross River | Obubra | ✓ | - | - | - | - | ✓ |
| Cross River | Abi | - | - | - | - | - | ✓ |
| Cross River | Bekwara | - | ✓ | - | - | - | ✓ |
| Ekiti | Ekiti South West | - | - | - | ✓ | - | - |
| Ekiti | Ikole | - | - | - | ✓ | - | - |
| Ekiti | Ekiti East | - | - | - | ✓ | - | - |
| Ekiti | Ijero | - | - | - | ✓ | - | - |
| Ekiti | Ilejemeje | - | - | - | ✓ | - | - |
| Ekiti | Efon Alaaye | - | - | - | ✓ | - | - |
| Ekiti | Irepodun /Ifelodun | - | - | - | ✓ | - | - |
| Ekiti | Ikere | - | - | - | ✓ | - | - |
| Ekiti | Ido Osi | - | - | - | ✓ | - | - |
| Ekiti | Ado-Ekiti | - | - | - | ✓ | - | - |
| Ekiti | Oye | - | - | - | ✓ | - | - |
| Ekiti | Gbonyin | - | - | - | - | - | ✓ |
| Ekiti | Emure | - | - | - | ✓ | - | - |
| Ekiti | Ekiti West | ✓ | - | - | ✓ | - | - |
| Ekiti | Ise Orun | ✓ | - | - | ✓ | - | - |
| Ekiti | Moba | ✓ | - | - | ✓ | - | - |
| FCT | Abuja Municipal Area Council | - | ✓ | - | ✓ | - | - |
| FCT | Kwali | - | ✓ | - | - | - | ✓ |
| FCT | Abaji | - | ✓ | - | - | - | ✓ |
| FCT | Bwari | ✓ | - | - | - | - | ✓ |
| Jigawa | Jahun | - | ✓ | - | - | - | ✓ |
| Jigawa | Garki | - | ✓ | - | - | - | ✓ |
| Jigawa | Auyo | - | ✓ | - | - | - | ✓ |
| Jigawa | Gwaram | - | ✓ | - | - | - | ✓ |
| Jigawa | Birnin Kudu | - | ✓ | - | - | - | ✓ |
| Jigawa | Kazaure | - | ✓ | - | - | - | ✓ |
| Jigawa | Miga | - | ✓ | - | - | - | ✓ |
| Jigawa | Kiri Kasamma | - | ✓ | - | - | - | ✓ |
| Jigawa | Sule Tankarkar | - | ✓ | - | - | - | ✓ |
| Jigawa | Gagarawa | - | ✓ | - | - | - | ✓ |
| Jigawa | Kaugama | - | ✓ | - | - | - | ✓ |
| Jigawa | Maigatari | - | ✓ | - | - | - | ✓ |
| Jigawa | Babura | - | ✓ | - | - | - | ✓ |
| Jigawa | Kiyawa | ✓ | - | - | - | - | ✓ |
| Jigawa | Ringim | ✓ | - | - | - | - | ✓ |
| Jigawa | Gwiwa | ✓ | - | - | - | - | ✓ |

| State | LGA Name | Schistosomiasis | | | STH | | |
|--------|--------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Jigawa | Kafin Hausa | ✓ | - | - | - | - | ✓ |
| Jigawa | Dutse | ✓ | - | - | - | - | ✓ |
| Jigawa | Taura | ✓ | - | - | - | - | ✓ |
| Jigawa | Buji | ✓ | - | - | - | - | ✓ |
| Jigawa | Roni | ✓ | - | - | - | - | ✓ |
| Jigawa | Yankwashi | ✓ | - | - | - | - | ✓ |
| Jigawa | Biriniwa | ✓ | - | - | - | - | ✓ |
| Jigawa | Gumel | ✓ | - | - | - | - | ✓ |
| Jigawa | Guri | ✓ | - | - | - | - | ✓ |
| Jigawa | Malam Madori | ✓ | - | - | - | - | ✓ |
| Jigawa | Hadejia | ✓ | - | - | - | - | - |
| Kaduna | Kudan | - | ✓ | - | ✓ | - | - |
| Kaduna | Sabon Gari | - | ✓ | - | ✓ | - | - |
| Kaduna | Soba | - | ✓ | - | ✓ | - | - |
| Kaduna | Giwa | - | ✓ | - | ✓ | - | - |
| Kaduna | Chikun | - | ✓ | - | ✓ | - | - |
| Kaduna | Sanga | - | ✓ | - | - | - | ✓ |
| Kaduna | Makarfi | - | ✓ | - | - | - | ✓ |
| Kaduna | Birnin Gwari | - | ✓ | - | ✓ | - | - |
| Kaduna | Zaria | - | ✓ | - | - | - | ✓ |
| Kaduna | Kauru | - | ✓ | - | ✓ | - | - |
| Kaduna | Ikara | - | ✓ | - | ✓ | - | - |
| Kaduna | Jema'a | - | ✓ | - | - | - | ✓ |
| Kaduna | Kachia | - | ✓ | - | - | - | ✓ |
| Kaduna | Kaura | - | ✓ | - | - | - | ✓ |
| Kaduna | Kaduna South | ✓ | - | - | ✓ | - | - |
| Kaduna | Jaba | ✓ | - | - | - | - | ✓ |
| Kaduna | Kubau | ✓ | - | - | ✓ | - | - |
| Kaduna | Kagarko | ✓ | - | - | - | - | ✓ |
| Kaduna | Igabi | ✓ | - | - | ✓ | - | - |
| Kaduna | Kajuru | ✓ | - | - | ✓ | - | - |
| Kaduna | Zangon Kataf | ✓ | - | - | ✓ | - | - |
| Kaduna | Lere | ✓ | - | - | ✓ | - | - |
| Kaduna | Kaduna North | ✓ | - | - | - | - | ✓ |
| Kano | Rimin Gado | - | ✓ | - | ✓ | - | - |
| Kano | Tudun Wada | - | ✓ | - | - | - | ✓ |
| Kano | Tofa | - | ✓ | - | ✓ | - | - |
| Kano | Kabo | - | ✓ | - | ✓ | - | - |
| Kano | Doguwa | - | ✓ | - | - | - | ✓ |
| Kano | Shanono | - | ✓ | - | ✓ | - | - |
| Kano | Madobi | - | ✓ | - | ✓ | - | - |
| Kano | Sumaila | - | ✓ | - | ✓ | - | - |
| Kano | Karaye | - | ✓ | - | ✓ | - | - |

| State | LGA Name | Schistosomiasis | | | STH | | |
|---------|----------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Kano | Minjibir | - | ✓ | - | - | - | ✓ |
| Kano | Kura | - | ✓ | - | ✓ | - | - |
| Kano | Kiru | - | ✓ | - | ✓ | - | - |
| Kano | Warawa | - | ✓ | - | ✓ | - | - |
| Kano | Rogo | - | ✓ | - | ✓ | - | - |
| Kano | Kunchi | - | ✓ | - | ✓ | - | - |
| Kano | Bagwai | - | ✓ | - | ✓ | - | - |
| Kano | Kibiya | - | ✓ | - | - | - | ✓ |
| Kano | Bunkure | - | ✓ | - | ✓ | - | - |
| Kano | Rano | - | ✓ | - | - | - | ✓ |
| Kano | Garko | - | ✓ | - | - | - | ✓ |
| Kano | Wudil | - | ✓ | - | - | - | ✓ |
| Kano | Garun Mallam | - | ✓ | - | - | - | ✓ |
| Kano | Takai | - | ✓ | - | - | - | ✓ |
| Kano | Gwarzo | - | ✓ | - | - | - | ✓ |
| Kano | Bebeji | - | ✓ | - | - | - | ✓ |
| Kano | Kano Municipal | ✓ | - | - | - | - | ✓ |
| Kano | Nasarawa | ✓ | - | - | - | - | ✓ |
| Kano | Dambatta | ✓ | - | - | - | - | ✓ |
| Kano | Dawakin Tofa | ✓ | - | - | - | - | ✓ |
| Kano | Dala | ✓ | - | - | - | - | ✓ |
| Kano | Gezawa | ✓ | - | - | ✓ | - | - |
| Kano | Bichi | ✓ | - | - | - | - | ✓ |
| Kano | Gwale | ✓ | - | - | - | - | ✓ |
| Kano | Makoda | ✓ | - | - | - | - | ✓ |
| Kano | Tarauni | ✓ | - | - | - | - | ✓ |
| Kano | Fagge | ✓ | - | - | - | - | ✓ |
| Kano | Ajingi | ✓ | - | - | - | - | ✓ |
| Kano | Dawakin Kudu | ✓ | - | - | - | - | ✓ |
| Kano | Gaya | ✓ | - | - | - | - | ✓ |
| Kano | Gabasawa | ✓ | - | - | ✓ | - | - |
| Kano | Tsanyawa | ✓ | - | - | ✓ | - | - |
| Kano | Kumbotso | ✓ | - | - | - | - | ✓ |
| Kano | Albasu | ✓ | - | - | - | - | ✓ |
| Kano | Ungogo | ✓ | - | - | - | - | ✓ |
| Katsina | Kafur | - | ✓ | - | - | - | ✓ |
| Katsina | Kusada | - | ✓ | - | - | - | ✓ |
| Katsina | Mani | - | ✓ | - | - | - | ✓ |
| Katsina | Dan Musa | - | ✓ | - | - | - | ✓ |
| Katsina | Matazu | - | ✓ | - | - | - | ✓ |
| Katsina | Kankia | - | ✓ | - | - | - | ✓ |
| Katsina | Musawa | - | ✓ | - | - | - | ✓ |
| Katsina | Dutsin Ma | - | ✓ | - | ✓ | - | - |

| State | LGA Name | Schistosomiasis | | | STH | | |
|---------|--------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Katsina | Faskari | - | ✓ | - | ✓ | - | - |
| Katsina | Safana | - | ✓ | - | - | - | ✓ |
| Katsina | Bindawa | - | ✓ | - | - | - | ✓ |
| Katsina | Bakori | - | ✓ | - | - | - | ✓ |
| Katsina | Kankara | - | ✓ | - | - | - | ✓ |
| Katsina | Dandume | - | ✓ | - | - | - | ✓ |
| Katsina | Rimi | - | ✓ | - | - | - | ✓ |
| Katsina | Kaita | - | ✓ | - | - | - | ✓ |
| Katsina | Malumfashi | - | ✓ | - | - | - | ✓ |
| Katsina | Katsina LGA | - | ✓ | - | - | - | ✓ |
| Katsina | Zango | - | - | - | - | - | ✓ |
| Katsina | Baure | - | - | - | - | - | ✓ |
| Katsina | Mashi | ✓ | - | - | - | - | ✓ |
| Katsina | Mai'Adua | ✓ | - | - | - | - | ✓ |
| Katsina | Ingawa | ✓ | - | - | ✓ | - | - |
| Katsina | Danja | ✓ | - | - | ✓ | - | - |
| Katsina | Funtua | ✓ | - | - | - | - | ✓ |
| Katsina | Charanchi | ✓ | - | - | - | - | ✓ |
| Katsina | Batsari | ✓ | - | - | - | - | ✓ |
| Katsina | Daura | ✓ | - | - | - | - | ✓ |
| Katsina | Sabuwa | ✓ | - | - | - | - | ✓ |
| Katsina | Kurfi | ✓ | - | - | - | - | ✓ |
| Katsina | Batagarawa | ✓ | - | - | - | - | ✓ |
| Katsina | Jibia | ✓ | - | - | - | - | ✓ |
| Katsina | Dutsi | ✓ | - | - | - | - | ✓ |
| Katsina | Sandamu | ✓ | - | - | - | - | ✓ |
| Kebbi | Koko/Bese | - | ✓ | - | - | - | ✓ |
| Kebbi | Ngaski | - | - | ✓ | - | - | ✓ |
| Kebbi | Danko/Wasagu | - | ✓ | - | - | - | ✓ |
| Kebbi | Shanga | - | ✓ | - | - | - | ✓ |
| Kebbi | Dandi | - | ✓ | - | ✓ | - | - |
| Kebbi | Bagudo | - | ✓ | - | ✓ | - | - |
| Kebbi | Yauri | - | ✓ | - | - | - | ✓ |
| Kebbi | Gwandu | - | ✓ | - | - | - | ✓ |
| Kebbi | Maiyama | - | ✓ | - | - | - | ✓ |
| Kebbi | Suru | - | ✓ | - | - | - | ✓ |
| Kebbi | Augie | - | - | ✓ | - | - | ✓ |
| Kebbi | Birnin Kebbi | - | ✓ | - | - | - | ✓ |
| Kebbi | Jega | - | ✓ | - | - | - | ✓ |
| Kebbi | Aleiro | - | ✓ | - | - | - | ✓ |
| Kebbi | Fakai | - | ✓ | - | - | - | ✓ |
| Kebbi | Sakaba | ✓ | - | - | - | - | ✓ |
| Kebbi | Bunza | ✓ | - | - | - | - | ✓ |

| State | LGA Name | Schistosomiasis | | | STH | | |
|-------|-----------------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Kebbi | Arewa | ✓ | - | - | - | - | ✓ |
| Kebbi | Zuru | ✓ | - | - | - | - | ✓ |
| Kebbi | Kalgo | ✓ | - | - | - | - | ✓ |
| Kogi | Mopa-Muro | - | - | - | ✓ | - | - |
| Kogi | Lokoja | - | ✓ | - | ✓ | - | - |
| Kogi | Omala | - | - | - | ✓ | - | - |
| Kogi | Ibaji | - | ✓ | - | ✓ | - | - |
| Kogi | Dekina | - | - | - | ✓ | - | - |
| Kogi | Kogi LGA | - | ✓ | - | ✓ | - | - |
| Kogi | Olamaboro | - | - | - | ✓ | - | - |
| Kogi | Ogori/Magongo | - | - | - | ✓ | - | - |
| Kogi | Yagba West | ✓ | - | - | - | - | ✓ |
| Kogi | Ankpa | ✓ | - | - | ✓ | - | - |
| Kogi | Idah | ✓ | - | - | ✓ | - | - |
| Kogi | Bassa | ✓ | - | - | ✓ | - | - |
| Kogi | Okehi | ✓ | - | - | ✓ | - | - |
| Kogi | Okene | ✓ | - | - | ✓ | - | - |
| Kogi | Ofu | ✓ | - | - | ✓ | - | - |
| Kogi | Ijumu | ✓ | - | - | ✓ | - | - |
| Kogi | Kabba/Bunu | ✓ | - | - | - | - | ✓ |
| Kogi | Yagba East | ✓ | - | - | - | - | ✓ |
| Kogi | Igalamela Odolu | ✓ | - | - | ✓ | - | - |
| Kogi | Ajaokuta | ✓ | - | - | ✓ | - | - |
| Kogi | Adavi | ✓ | - | - | ✓ | - | - |
| Lagos | Ojo | - | - | - | - | ✓ | - |
| Lagos | Apapa | - | - | - | ✓ | - | - |
| Lagos | Badagry | - | - | - | ✓ | - | - |
| Lagos | Lagos Island | - | - | - | ✓ | - | - |
| Lagos | Shomolu | - | - | - | ✓ | - | - |
| Lagos | Mushin | - | - | - | - | - | ✓ |
| Lagos | Ibeju-Lekki | ✓ | - | - | - | ✓ | - |
| Lagos | Amuwo-Odofin | ✓ | - | - | - | ✓ | - |
| Lagos | Eti Osa | ✓ | - | - | ✓ | - | - |
| Lagos | Ajeromi-Ifelodun | ✓ | - | - | ✓ | - | - |
| Lagos | Lagos Mainland | ✓ | - | - | ✓ | - | - |
| Lagos | Surulere, Lagos State | ✓ | - | - | - | - | ✓ |
| Lagos | Ifako-Ijaiye | ✓ | - | - | - | - | ✓ |
| Lagos | Agege | ✓ | - | - | - | - | ✓ |
| Lagos | Kosofe | ✓ | - | - | ✓ | - | - |
| Lagos | Ikorodu | ✓ | - | - | - | - | ✓ |
| Lagos | Ikeja | ✓ | - | - | - | - | ✓ |
| Lagos | Oshodi-Isolo | ✓ | - | - | - | - | ✓ |
| Lagos | Alimosho | ✓ | - | - | - | - | ✓ |

| State | LGA Name | Schistosomiasis | | | STH | | |
|-------|-----------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Niger | Chanchaga | - | ✓ | - | ✓ | - | - |
| Niger | Munya | - | ✓ | - | - | ✓ | - |
| Niger | Tafa | - | ✓ | - | - | - | ✓ |
| Niger | Bosso | - | ✓ | - | ✓ | - | - |
| Niger | Rijau | - | - | ✓ | ✓ | - | - |
| Niger | Rafi | - | ✓ | - | - | - | ✓ |
| Niger | Magama | - | ✓ | - | ✓ | - | - |
| Niger | Borgu | - | - | ✓ | - | - | ✓ |
| Niger | Agaie | - | ✓ | - | ✓ | - | - |
| Niger | Wushishi | - | ✓ | - | ✓ | - | - |
| Niger | Edati | - | ✓ | - | ✓ | - | - |
| Niger | Katcha | - | ✓ | - | ✓ | - | - |
| Niger | Mashegu | - | ✓ | - | ✓ | - | - |
| Niger | Mokwa | - | ✓ | - | ✓ | - | - |
| Niger | Gurara | - | ✓ | - | ✓ | - | - |
| Niger | Agwara | - | ✓ | - | ✓ | - | - |
| Niger | Bida | - | ✓ | - | ✓ | - | - |
| Niger | Mariga | - | ✓ | - | ✓ | - | - |
| Niger | Shiroro | - | ✓ | - | ✓ | - | - |
| Niger | Lavun | - | ✓ | - | ✓ | - | - |
| Niger | Suleja | - | ✓ | - | ✓ | - | - |
| Niger | Paikoro | - | ✓ | - | ✓ | - | - |
| Niger | Kontogora | - | ✓ | - | ✓ | - | - |
| Niger | Gbako | ✓ | - | - | - | ✓ | - |
| Niger | Lapai | ✓ | - | - | - | ✓ | - |
| Osun | Atakunmosa East | - | ✓ | - | - | ✓ | - |
| Osun | Ife North | - | ✓ | - | - | ✓ | - |
| Osun | Ejigbo | - | ✓ | - | ✓ | - | - |
| Osun | Boluwaduro | - | - | - | ✓ | - | - |
| Osun | Olorunda | - | ✓ | - | ✓ | - | - |
| Osun | Ifedayo | - | - | - | ✓ | - | - |
| Osun | Irewole | ✓ | - | - | - | ✓ | - |
| Osun | Osogbo | ✓ | - | - | ✓ | - | - |
| Osun | Oriade | ✓ | - | - | ✓ | - | - |
| Osun | Ife Central | ✓ | - | - | ✓ | - | - |
| Osun | Ilesha West | ✓ | - | - | ✓ | - | - |
| Osun | Aiyedaade | ✓ | - | - | ✓ | - | - |
| Osun | Ifelodun | ✓ | - | - | ✓ | - | - |
| Osun | Ife South | ✓ | - | - | - | ✓ | - |
| Osun | Ilesa East | ✓ | - | - | ✓ | - | - |
| Osun | Aiyedire | ✓ | - | - | ✓ | - | - |
| Osun | Obokun | ✓ | - | - | ✓ | - | - |
| Osun | Isokan | ✓ | - | - | - | ✓ | - |

| State | LGA Name | Schistosomiasis | | | STH | | |
|-------|-------------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Osun | Odo Otin | ✓ | - | - | ✓ | - | - |
| Osun | Ola Oluwa | ✓ | - | - | ✓ | - | - |
| Osun | Boripe | ✓ | - | - | ✓ | - | - |
| Osun | Atakunmosa West | ✓ | - | - | - | ✓ | - |
| Osun | Iwo | ✓ | - | - | - | ✓ | - |
| Osun | Egbedore | ✓ | - | - | ✓ | - | - |
| Osun | Ede South | ✓ | - | - | - | ✓ | - |
| Osun | Ife East | ✓ | - | - | - | ✓ | - |
| Osun | Ila | ✓ | - | - | ✓ | - | - |
| Osun | Ede North | ✓ | - | - | ✓ | - | - |
| Osun | Irepodun | ✓ | - | - | ✓ | - | - |
| Osun | Orolu | ✓ | - | - | - | ✓ | - |
| Oyo | Ibadan North West | - | - | - | ✓ | - | - |
| Oyo | Ido | - | - | - | ✓ | - | - |
| Oyo | Iwajowa | - | ✓ | - | ✓ | - | - |
| Oyo | Ogo Oluwa | - | ✓ | - | - | ✓ | - |
| Oyo | Saki West | - | ✓ | - | - | ✓ | - |
| Oyo | Saki East | - | ✓ | - | ✓ | - | - |
| Oyo | Ibadan North East | ✓ | - | - | ✓ | - | - |
| Oyo | Ibadan South East | ✓ | - | - | ✓ | - | - |
| Oyo | Ibadan South West | ✓ | - | - | ✓ | - | - |
| Oyo | Ibadan North | ✓ | - | - | ✓ | - | - |
| Oyo | Lagelu | ✓ | - | - | - | ✓ | - |
| Oyo | Akinyele | ✓ | - | - | - | ✓ | - |
| Oyo | Oyo West | ✓ | - | - | - | ✓ | - |
| Oyo | Afijio | ✓ | - | - | - | ✓ | - |
| Oyo | Oluyole | ✓ | - | - | - | ✓ | - |
| Oyo | Kajola | ✓ | - | - | ✓ | - | - |
| Oyo | Ona Ara | ✓ | - | - | - | ✓ | - |
| Oyo | Itesiwaju | ✓ | - | - | ✓ | - | - |
| Oyo | Atisbo | ✓ | - | - | - | - | ✓ |
| Oyo | Ogbomosho North | ✓ | - | - | ✓ | - | - |
| Oyo | Atiba | ✓ | - | - | - | ✓ | - |
| Oyo | Olorunsogo | ✓ | - | - | ✓ | - | - |
| Oyo | Ibarapa Central | ✓ | - | - | ✓ | - | - |
| Oyo | Ogbomosho South | ✓ | - | - | ✓ | - | - |
| Oyo | Ibarapa East | ✓ | - | - | ✓ | - | - |
| Oyo | Ori Ire | ✓ | - | - | ✓ | - | - |
| Oyo | Surulere, Oyo | ✓ | - | - | ✓ | - | - |
| Oyo | Ibarapa North | ✓ | - | - | ✓ | - | - |
| Oyo | Irepo | ✓ | - | - | - | ✓ | - |
| Oyo | Orelope | ✓ | - | - | ✓ | - | - |
| Oyo | Oyo East | ✓ | - | - | - | ✓ | - |

| State | LGA Name | Schistosomiasis | | | STH | | |
|--------|-------------------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Oyo | Iseyin | ✓ | - | - | - | ✓ | - |
| Oyo | Egbeda | ✓ | - | - | - | ✓ | - |
| Rivers | Abua/Odual | - | - | - | - | ✓ | - |
| Rivers | KHANA | - | - | - | - | ✓ | - |
| Rivers | Okrika | - | - | - | ✓ | - | - |
| Rivers | Andoni | - | - | - | - | ✓ | - |
| Rivers | Opobo/Nkoro | - | - | - | - | ✓ | - |
| Rivers | AHOADA WEST | - | - | - | ✓ | - | - |
| Rivers | Ogba/Egbema/Ndoni | - | - | - | ✓ | - | - |
| Rivers | Emohua | - | - | - | - | ✓ | - |
| Rivers | GOKANA | - | - | - | - | ✓ | - |
| Rivers | BONNY | - | - | - | - | ✓ | - |
| Rivers | Akuku Toru | - | - | - | ✓ | - | - |
| Rivers | Ikwerre | - | - | - | ✓ | - | - |
| Rivers | Ogu Bolo | - | - | - | ✓ | - | - |
| Rivers | Tai | - | - | - | ✓ | - | - |
| Rivers | Etche | - | - | - | ✓ | - | - |
| Rivers | Omuma | - | - | - | - | ✓ | - |
| Rivers | OYIGBO | - | - | - | - | - | ✓ |
| Rivers | AHOADA EAST | ✓ | - | - | - | ✓ | - |
| Rivers | DEGEMA | ✓ | - | - | ✓ | - | - |
| Rivers | Asari Toru | ✓ | - | - | ✓ | - | - |
| Rivers | Port Harcourt | ✓ | - | - | - | - | ✓ |
| Rivers | ELEME | ✓ | - | - | ✓ | - | - |
| Rivers | Obio/Akpor | ✓ | - | - | ✓ | - | - |
| Taraba | Gassol | - | ✓ | - | - | - | ✓ |
| Taraba | Bali | ✓ | - | - | - | - | ✓ |
| Taraba | Gashaka | ✓ | - | - | - | - | ✓ |
| Taraba | Donga | ✓ | - | - | - | - | ✓ |
| Taraba | Lau | ✓ | - | - | - | - | ✓ |
| Taraba | Zing | ✓ | - | - | - | - | ✓ |
| Taraba | Karim-Lamido | ✓ | - | - | - | - | ✓ |
| Taraba | Yorro | ✓ | - | - | - | - | ✓ |
| Yobe | Karasuwa | - | ✓ | - | - | - | ✓ |
| Yobe | Bursari | - | ✓ | - | - | - | ✓ |
| Yobe | Nguru | - | ✓ | - | - | - | ✓ |
| Yobe | Geidam | - | ✓ | - | - | - | ✓ |
| Yobe | Jakusko | - | ✓ | - | - | - | ✓ |
| Yobe | Yunusari | - | ✓ | - | - | - | ✓ |
| Yobe | Damaturu | - | ✓ | - | - | - | ✓ |
| Yobe | Bade | - | ✓ | - | - | - | ✓ |
| Yobe | Tarmuwa | - | ✓ | - | - | - | - |
| Yobe | Fune | ✓ | - | - | - | - | ✓ |

| State | LGA Name | Schistosomiasis | | | STH | | |
|-------|----------|-----------------|----------|------|-----|------|----------|
| | | Low | Moderate | High | Low | High | Case Mgt |
| Yobe | Nangere | ✓ | - | - | - | - | ✓ |
| Yobe | Fika | ✓ | - | - | - | - | ✓ |
| Yobe | Yusufari | ✓ | - | - | - | - | ✓ |
| Yobe | Potiskum | ✓ | - | - | - | - | ✓ |
| Yobe | Machina | ✓ | - | - | - | - | - |

Appendix 8: Schistosomiasis and STHs Prevalence by Practice of Defecation

| State | Defecation Practices | No of Respondents | Schistosomiasis | | STHs | |
|-------------|----------------------------|-------------------|-----------------|-------|-------------|--------|
| | | | No Infected | (%) | No Infected | (%) |
| Akwalbom | In the School Toilet | 4864 | 14 | .3% | 2621 | 53.9% |
| | Around school Compound | 1430 | 5 | .3% | 929 | 65.0% |
| | Outside school environment | 1514 | 3 | .3% | 1008 | 133.6% |
| | Others | 58 | 0 | 0.0% | 32 | 112.0% |
| | Total | 7866 | 22 | .3% | 4590 | 58.4% |
| Bauchi | In the School Toilet | 2099 | 185 | 8.8% | 149 | 7.1% |
| | Around school Compound | 1266 | 258 | 20.4% | 158 | 12.5% |
| | Outside school environment | 1535 | 231 | 15.2% | 187 | 19.9% |
| | Others | 58 | 1 | 1.8% | 2 | 3.5% |
| | Total | 4958 | 675 | 13.6% | 496 | 10.0% |
| Bayelsa | In the School Toilet | 856 | 5 | .6% | 218 | 25.5% |
| | Around school Compound | 386 | 4 | 1.0% | 153 | 39.6% |
| | Outside school environment | 548 | 5 | 2.3% | 210 | 75.0% |
| | Others | 153 | 3 | 2.0% | 58 | 37.9% |
| | Total | 1943 | 17 | .9% | 639 | 32.9% |
| Benue | In the School Toilet | 1455 | 214 | 14.7% | 431 | 29.6% |
| | Around school Compound | 802 | 90 | 11.2% | 191 | 23.8% |
| | Outside school environment | 1195 | 147 | 20.3% | 250 | 32.3% |
| | Others | 0 | 0 | 0.0% | 0 | 0.0% |
| | Total | 3452 | 451 | 13.1% | 872 | 25.3% |
| Cross River | In the School Toilet | 3058 | 178 | 5.8% | 830 | 27.1% |
| | Around school Compound | 1187 | 74 | 6.2% | 232 | 19.5% |
| | Outside school environment | 635 | 31 | 7.2% | 137 | 31.1% |
| | Others | 63 | 0 | 0.0% | 10 | 27.8% |
| | Total | 4943 | 283 | 5.7% | 1209 | 24.5% |
| Ekiti | In the School Toilet | 1139 | 1 | .1% | 317 | 27.8% |
| | Around school Compound | 1396 | 2 | .1% | 464 | 33.2% |
| | Outside school environment | 658 | 1 | .2% | 190 | 58.5% |
| | Others | 330 | 4 | 1.9% | 113 | 67.4% |
| | Total | 3523 | 8 | .2% | 1084 | 30.8% |
| FCT | In the School Toilet | 421 | 76 | 18.1% | 78 | 18.5% |
| | Around school Compound | 376 | 94 | 25.0% | 65 | 17.3% |
| | Outside school environment | 173 | 33 | 40.3% | 43 | 46.6% |
| | Others | 33 | 1 | 3.0% | 7 | 21.2% |
| | Total | 1003 | 204 | 20.3% | 193 | 19.2% |
| Jigawa | In the School Toilet | 2877 | 300 | 10.4% | 193 | 6.7% |
| | Around school Compound | 1222 | 132 | 10.8% | 59 | 4.8% |

| State | Defecation Practices | No of Respondents | Schistosomiasis | | STHs | |
|---------|----------------------------|-------------------|-----------------|-------|-------------|--------|
| | | | No Infected | (%) | No Infected | (%) |
| | Outside school environment | 2421 | 311 | 21.4% | 151 | 12.7% |
| | Others | 9 | 0 | 0.0% | 1 | 14.3% |
| | Total | 6529 | 743 | 11.4% | 404 | 6.2% |
| | | | | | | |
| Kaduna | In the School Toilet | 1834 | 209 | 11.4% | 368 | 20.1% |
| | Around school Compound | 1274 | 184 | 14.4% | 246 | 19.3% |
| | Outside school environment | 2652 | 401 | 29.6% | 643 | 47.0% |
| | Others | 101 | 17 | 32.2% | 22 | 41.2% |
| | Total | 5861 | 811 | 13.8% | 1279 | 21.8% |
| Kano | In the School Toilet | 6273 | 730 | 11.6% | 957 | 15.3% |
| | Around school Compound | 1128 | 162 | 14.4% | 184 | 16.3% |
| | Outside school environment | 3394 | 581 | 36.5% | 722 | 42.9% |
| | Others | 209 | 58 | 45.9% | 60 | 43.3% |
| | Total | 11004 | 1531 | 13.9% | 1923 | 17.5% |
| Katsina | In the School Toilet | 4390 | 395 | 9.0% | 381 | 8.7% |
| | Around school Compound | 1871 | 288 | 15.4% | 238 | 12.7% |
| | Outside school environment | 2021 | 261 | 23.2% | 251 | 23.5% |
| | Others | 54 | 0 | 0.0% | 2 | 3.8% |
| | Total | 8336 | 944 | 11.3% | 872 | 10.5% |
| Kebbi | In the School Toilet | 1382 | 244 | 17.7% | 98 | 7.1% |
| | Around school Compound | 1851 | 435 | 23.5% | 206 | 11.1% |
| | Outside school environment | 1611 | 381 | 42.4% | 175 | 21.7% |
| | Others | 10 | 2 | 62.5% | 1 | 12.5% |
| | Total | 4854 | 1062 | 21.9% | 480 | 9.9% |
| Kogi | In the School Toilet | 896 | 31 | 3.5% | 215 | 24.0% |
| | Around school Compound | 1019 | 27 | 2.6% | 263 | 25.8% |
| | Outside school environment | 2569 | 77 | 5.5% | 789 | 63.1% |
| | Others | 788 | 14 | 2.0% | 214 | 56.3% |
| | Total | 5272 | 149 | 2.8% | 1481 | 28.1% |
| Lagos | In the School Toilet | 3767 | 33 | .9% | 890 | 23.6% |
| | Around school Compound | 740 | 7 | .9% | 316 | 42.7% |
| | Outside school environment | 209 | 1 | .9% | 102 | 96.9% |
| | Others | 58 | 0 | 0.0% | 33 | 155.4% |
| | Total | 4774 | 41 | .9% | 1341 | 28.1% |
| Niger | In the School Toilet | 1531 | 495 | 32.3% | 582 | 38.0% |
| | Around school Compound | 3719 | 938 | 25.2% | 1376 | 37.0% |
| | Outside school environment | 1788 | 406 | 49.5% | 498 | 58.3% |
| | Others | 159 | 40 | 44.2% | 75 | 86.8% |
| | Total | 7197 | 1879 | 26.1% | 2531 | 35.2% |
| Osun | In the School Toilet | 2445 | 109 | 4.5% | 988 | 40.4% |

| State | Defecation Practices | No of Respondents | Schistosomiasis | | STHs | |
|--------|----------------------------|-------------------|-----------------|-------|-------------|--------|
| | | | No Infected | (%) | No Infected | (%) |
| | Around school Compound | 2456 | 151 | 6.1% | 1123 | 45.7% |
| | Outside school environment | 2606 | 145 | 11.5% | 1284 | 118.9% |
| | Others | 72 | 0 | 0.0% | 31 | 43.1% |
| | Total | 7579 | 405 | 5.3% | 3426 | 45.2% |
| | | | | | | |
| Oyo | In the School Toilet | 1808 | 94 | 5.2% | 839 | 46.4% |
| | Around school Compound | 3969 | 234 | 5.9% | 2023 | 51.0% |
| | Outside school environment | 2332 | 107 | 10.5% | 966 | 89.3% |
| | Others | 1 | 0 | 0.0% | 0 | 0.0% |
| | Total | 8110 | 435 | 5.4% | 3828 | 47.2% |
| Rivers | In the School Toilet | 2871 | 5 | .2% | 1113 | 38.8% |
| | Around school Compound | 2048 | 2 | .1% | 994 | 48.5% |
| | Outside school environment | 774 | 0 | 0.0% | 351 | 79.6% |
| | Others | 27 | 0 | 0.0% | 9 | 33.3% |
| | Total | 5720 | 7 | .1% | 2467 | 43.1% |
| Taraba | In the School Toilet | 409 | 17 | 4.2% | 17 | 4.2% |
| | Around school Compound | 433 | 31 | 7.2% | 20 | 4.6% |
| | Outside school environment | 981 | 53 | 7.7% | 62 | 8.7% |
| | Others | 24 | 2 | 20.0% | 4 | 34.3% |
| | Total | 1847 | 103 | 5.6% | 103 | 5.6% |
| Yobe | In the School Toilet | 1351 | 166 | 12.3% | 29 | 2.1% |
| | Around school Compound | 1274 | 201 | 15.8% | 10 | .8% |
| | Outside school environment | 976 | 195 | 34.6% | 9 | 1.1% |
| | Others | 100 | 17 | 17.0% | 3 | 3.0% |
| | Total | 3701 | 579 | 15.6% | 51 | 1.4% |

Appendix 9: Prevalence of Schistosomiasis and STHs by water contact activities


| State | Water Contact Activities | No of Respondents | Schistosomiasis | | STHs | |
|-----------|--------------------------|-------------------|-----------------|-------|-------------|-------|
| | | | No Infected | (%) | No Infected | (%) |
| Akwa Ibom | Bathing | 2680 | 4 | .1% | 1608 | 60.0% |
| | Washing | 1423 | 4 | .3% | 828 | 58.2% |
| | Fishing | 81 | 3 | 3.7% | 51 | 63.0% |
| | Crossing Water | 68 | 1 | 1.5% | 42 | 61.8% |
| | Fetching Water | 1824 | 6 | .3% | 1129 | 61.9% |
| | Playing | 495 | 2 | .4% | 278 | 56.2% |
| | Swimming | 355 | 0 | 0.0% | 222 | 62.5% |
| | Others | 940 | 2 | .2% | 432 | 46.0% |
| | Total | 7866 | 22 | .3% | 4590 | 58.4% |
| Bauchi | Bathing | 509 | 97 | 19.1% | 61 | 12.0% |
| | Washing | 600 | 126 | 21.0% | 53 | 8.8% |
| | Fishing | 18 | 6 | 33.3% | 3 | 16.7% |
| | Crossing Water | 113 | 18 | 15.9% | 9 | 8.0% |
| | Fetching Water | 462 | 87 | 18.8% | 36 | 7.8% |
| | Playing | 1240 | 125 | 10.1% | 157 | 12.7% |
| | Swimming | 115 | 22 | 19.1% | 12 | 10.4% |
| | Others | 1901 | 194 | 10.2% | 165 | 8.7% |
| | Total | 4958 | 675 | 13.6% | 496 | 10.0% |
| Bayelsa | Bathing | 254 | 3 | 1.2% | 77 | 30.3% |
| | Washing | 148 | 3 | 2.0% | 64 | 43.2% |
| | Fishing | 19 | 0 | 0.0% | 9 | 47.4% |
| | Crossing Water | 19 | 1 | 5.3% | 13 | 68.4% |
| | Fetching Water | 594 | 2 | .3% | 178 | 30.0% |
| | Playing | 219 | 4 | 1.8% | 76 | 34.7% |
| | Swimming | 527 | 3 | .6% | 188 | 35.7% |
| | Others | 163 | 1 | .6% | 34 | 20.9% |
| | Total | 1943 | 17 | .9% | 639 | 32.9% |
| Benue | Bathing | 685 | 92 | 13.4% | 228 | 33.3% |
| | Washing | 856 | 116 | 13.6% | 212 | 24.8% |
| | Fishing | 203 | 28 | 13.8% | 47 | 23.2% |
| | Crossing Water | 132 | 20 | 15.2% | 37 | 28.0% |
| | Fetching Water | 289 | 38 | 13.1% | 79 | 27.3% |
| | Playing | 904 | 98 | 10.8% | 193 | 21.3% |
| | Swimming | 210 | 32 | 15.2% | 41 | 19.5% |
| | Others | 173 | 27 | 15.6% | 35 | 20.2% |
| | Total | 3452 | 451 | 13.1% | 872 | 25.3% |

| State | Water Contact Activities | No of Respondents | Schistosomiasis | | STHs | |
|-------------|--------------------------|-------------------|-----------------|-------|-------------|-------|
| | | | No Infected | (%) | No Infected | (%) |
| Cross River | Bathing | 1503 | 80 | 5.3% | 393 | 26.1% |
| | Washing | 928 | 56 | 6.0% | 177 | 19.1% |
| | Fishing | 138 | 10 | 7.2% | 35 | 25.4% |
| | Crossing Water | 131 | 10 | 7.6% | 16 | 12.2% |
| | Fetching Water | 1434 | 92 | 6.4% | 385 | 26.8% |
| | Playing | 160 | 2 | 1.3% | 43 | 26.9% |
| | Swimming | 206 | 27 | 13.1% | 47 | 22.8% |
| | Others | 443 | 6 | 1.4% | 113 | 25.5% |
| | Total | 4943 | 283 | 5.7% | 1209 | 24.5% |
| Ekiti | Bathing | 480 | 0 | 0.0% | 146 | 30.4% |
| | Washing | 732 | 0 | 0.0% | 182 | 24.9% |
| | Fishing | 46 | 0 | 0.0% | 10 | 21.7% |
| | Crossing Water | 79 | 0 | 0.0% | 12 | 15.2% |
| | Fetching Water | 569 | 1 | .2% | 169 | 29.7% |
| | Playing | 336 | 0 | 0.0% | 101 | 30.1% |
| | Swimming | 85 | 0 | 0.0% | 23 | 27.1% |
| | Others | 1196 | 7 | .6% | 441 | 36.9% |
| | Total | 3523 | 8 | .2% | 1084 | 30.8% |
| FCT | Bathing | 217 | 55 | 25.3% | 43 | 19.8% |
| | Washing | 174 | 27 | 15.5% | 41 | 23.6% |
| | Fishing | 13 | 2 | 15.4% | 4 | 30.8% |
| | Crossing Water | 7 | 3 | 42.9% | 0 | 0.0% |
| | Fetching Water | 142 | 18 | 12.7% | 30 | 21.1% |
| | Playing | 140 | 36 | 25.7% | 22 | 15.7% |
| | Swimming | 96 | 26 | 27.1% | 22 | 22.9% |
| | Others | 214 | 37 | 17.3% | 31 | 14.5% |
| | Total | 1003 | 204 | 20.3% | 193 | 19.2% |
| Jigawa | Bathing | 790 | 70 | 8.9% | 55 | 7.0% |
| | Washing | 1327 | 150 | 11.3% | 106 | 8.0% |
| | Fishing | 151 | 24 | 15.9% | 13 | 8.6% |
| | Crossing Water | 571 | 75 | 13.1% | 36 | 6.3% |
| | Fetching Water | 853 | 122 | 14.3% | 58 | 6.8% |
| | Playing | 1426 | 135 | 9.5% | 55 | 3.9% |
| | Swimming | 263 | 58 | 22.1% | 15 | 5.7% |
| | Others | 1148 | 109 | 9.5% | 66 | 5.7% |
| | Total | 6529 | 743 | 11.4% | 404 | 6.2% |
| Kaduna | Bathing | 755 | 128 | 17.0% | 207 | 27.4% |
| | Washing | 842 | 95 | 11.3% | 158 | 18.8% |

| State | Water Contact Activities | No of Respondents | Schistosomiasis | | STHs | |
|----------------|--------------------------|-------------------|-----------------|-------|-------------|-------|
| | | | No Infected | (%) | No Infected | (%) |
| | Fishing | 50 | 6 | 12.0% | 12 | 24.0% |
| | Crossing Water | 231 | 18 | 7.8% | 30 | 13.0% |
| | Fetching Water | 669 | 67 | 10.0% | 147 | 22.0% |
| | Playing | 1585 | 247 | 15.6% | 351 | 22.1% |
| | Swimming | 690 | 134 | 19.4% | 167 | 24.2% |
| | Others | 1039 | 116 | 11.2% | 207 | 19.9% |
| | Total | 5861 | 811 | 13.8% | 1279 | 21.8% |
| | Kano | Bathing | 203 | 104 | 51.2% | 41 |
| Washing | | 747 | 204 | 27.3% | 154 | 20.6% |
| Fishing | | 149 | 68 | 45.6% | 22 | 14.8% |
| Crossing Water | | 42 | 10 | 23.8% | 8 | 19.0% |
| Fetching Water | | 360 | 34 | 9.4% | 43 | 11.9% |
| Playing | | 889 | 161 | 18.1% | 172 | 19.3% |
| Swimming | | 337 | 94 | 27.9% | 59 | 17.5% |
| Total | | 11004 | 1531 | 13.9% | 1923 | 17.5% |
| Katsina | Bathing | 705 | 131 | 18.6% | 104 | 14.8% |
| | Washing | 2374 | 255 | 10.7% | 212 | 8.9% |
| | Fishing | 7 | 1 | 14.3% | 0 | 0.0% |
| | Crossing Water | 57 | 7 | 12.3% | 5 | 8.8% |
| | Fetching Water | 812 | 93 | 11.5% | 72 | 8.9% |
| | Playing | 211 | 35 | 16.6% | 21 | 10.0% |
| | Swimming | 250 | 76 | 30.4% | 45 | 18.0% |
| | Total | 8336 | 944 | 11.3% | 872 | 10.5% |
| Kebbi | Bathing | 689 | 157 | 22.8% | 63 | 9.1% |
| | Washing | 422 | 90 | 21.3% | 36 | 8.5% |
| | Fishing | 228 | 71 | 31.1% | 44 | 19.3% |
| | Crossing Water | 144 | 41 | 28.5% | 16 | 11.1% |
| | Fetching Water | 529 | 110 | 20.8% | 51 | 9.6% |
| | Playing | 1292 | 311 | 24.1% | 111 | 8.6% |
| | Swimming | 740 | 194 | 26.2% | 89 | 12.0% |
| | Total | 4854 | 1062 | 21.9% | 480 | 9.9% |
| Kogi | Bathing | 574 | 18 | 3.1% | 166 | 28.9% |
| | Washing | 704 | 16 | 2.3% | 244 | 34.7% |
| | Fishing | 51 | 1 | 2.0% | 13 | 25.5% |
| | Crossing Water | 111 | 4 | 3.6% | 40 | 36.0% |

| State | Water Contact Activities | No of Respondents | Schistosomiasis | | STHs | |
|-------|--------------------------|-------------------|-----------------|-------|-------------|-------|
| | | | No Infected | (%) | No Infected | (%) |
| | Fetching Water | 944 | 50 | 5.3% | 250 | 26.5% |
| | Playing | 1024 | 35 | 3.4% | 240 | 23.4% |
| | Swimming | 326 | 13 | 4.0% | 103 | 31.6% |
| | Others | 1538 | 12 | .8% | 425 | 27.6% |
| | Total | 5272 | 149 | 2.8% | 1481 | 28.1% |
| Lagos | Bathing | 85 | 1 | 1.2% | 64 | 75.3% |
| | Washing | 758 | 3 | .4% | 193 | 25.5% |
| | Fishing | 28 | 1 | 3.6% | 21 | 75.0% |
| | Crossing Water | 45 | 0 | 0.0% | 24 | 53.3% |
| | Fetching Water | 161 | 4 | 2.5% | 79 | 49.1% |
| | Playing | 69 | 0 | 0.0% | 45 | 65.2% |
| | Swimming | 117 | 0 | 0.0% | 78 | 66.7% |
| | Others | 3511 | 32 | .9% | 837 | 23.8% |
| | Total | 4774 | 41 | .9% | 1341 | 28.1% |
| Niger | Bathing | 899 | 273 | 30.4% | 415 | 46.2% |
| | Washing | 2552 | 643 | 25.2% | 890 | 34.9% |
| | Fishing | 232 | 55 | 23.7% | 62 | 26.7% |
| | Crossing Water | 188 | 27 | 14.4% | 77 | 41.0% |
| | Fetching Water | 940 | 227 | 24.1% | 367 | 39.0% |
| | Playing | 628 | 179 | 28.5% | 148 | 23.6% |
| | Swimming | 862 | 328 | 38.1% | 241 | 28.0% |
| | Others | 896 | 147 | 16.4% | 331 | 36.9% |
| | Total | 7197 | 1879 | 26.1% | 2531 | 35.2% |
| Osun | Bathing | 893 | 36 | 4.0% | 408 | 45.7% |
| | Washing | 2624 | 175 | 6.7% | 1296 | 49.4% |
| | Fishing | 79 | 27 | 34.2% | 35 | 44.3% |
| | Crossing Water | 140 | 5 | 3.6% | 70 | 50.0% |
| | Fetching Water | 1349 | 55 | 4.1% | 626 | 46.4% |
| | Playing | 494 | 20 | 4.0% | 213 | 43.1% |
| | Swimming | 233 | 11 | 4.7% | 126 | 54.1% |
| | Others | 1767 | 76 | 4.3% | 652 | 36.9% |
| | Total | 7579 | 405 | 5.3% | 3426 | 45.2% |
| Oyo | Bathing | 566 | 26 | 4.6% | 254 | 44.9% |
| | Washing | 1506 | 44 | 2.9% | 648 | 43.0% |
| | Fishing | 145 | 10 | 6.9% | 67 | 46.2% |
| | Crossing Water | 451 | 30 | 6.7% | 228 | 50.6% |
| | Fetching Water | 1783 | 71 | 4.0% | 690 | 38.7% |
| | Playing | 1710 | 58 | 3.4% | 808 | 47.3% |

| State | Water Contact Activities | No of Respondents | Schistosomiasis | | STHs | |
|--------|--------------------------|-------------------|-----------------|-------|-------------|-------|
| | | | No Infected | (%) | No Infected | (%) |
| | Swimming | 825 | 73 | 8.8% | 472 | 57.2% |
| | Others | 1124 | 123 | 10.9% | 661 | 58.8% |
| | Total | 8110 | 435 | 5.4% | 3828 | 47.2% |
| | | | | | | |
| Rivers | Bathing | 914 | 2 | .2% | 394 | 43.1% |
| | Washing | 1359 | 2 | .1% | 659 | 48.5% |
| | Fishing | 1714 | 3 | .2% | 750 | 43.8% |
| | Crossing Water | 616 | 0 | 0.0% | 252 | 40.9% |
| | Fetching Water | 116 | 0 | 0.0% | 49 | 42.2% |
| | Playing | 343 | 0 | 0.0% | 153 | 44.6% |
| | Swimming | 9 | 0 | 0.0% | 3 | 33.3% |
| | Others | 649 | 0 | 0.0% | 207 | 31.9% |
| | Total | 5720 | 7 | .1% | 2467 | 43.1% |
| Taraba | Bathing | 112 | 6 | 5.4% | 7 | 6.3% |
| | Washing | 54 | 5 | 9.3% | 3 | 5.6% |
| | Fishing | 15 | 4 | 26.7% | 1 | 6.7% |
| | Crossing Water | 15 | 0 | 0.0% | 0 | 0.0% |
| | Fetching Water | 943 | 42 | 4.5% | 42 | 4.5% |
| | Playing | 76 | 4 | 5.3% | 12 | 15.8% |
| | Swimming | 169 | 4 | 2.4% | 17 | 10.1% |
| | Others | 463 | 38 | 8.2% | 21 | 4.5% |
| | Total | 1847 | 103 | 5.6% | 103 | 5.6% |
| Yobe | Bathing | 322 | 68 | 21.1% | 5 | 1.6% |
| | Washing | 559 | 92 | 16.5% | 4 | .7% |
| | Fishing | 168 | 29 | 17.3% | 0 | 0.0% |
| | Crossing Water | 93 | 9 | 9.7% | 2 | 2.2% |
| | Fetching Water | 31 | 5 | 16.1% | 1 | 3.2% |
| | Playing | 1123 | 172 | 15.3% | 24 | 2.1% |
| | Swimming | 175 | 24 | 13.7% | 0 | 0.0% |
| | Others | 1230 | 180 | 14.6% | 15 | 1.2% |
| | Total | 3701 | 579 | 15.6% | 51 | 1.4% |



Legend

- 0%
- <10% but >0%
- ≥10% but <50%
- ≥50%

