



PSI Research & Metrics Quarterly Scorecard

Indicator	Q1 2007	Q2 2007	Q3 2007	Q4 2007	Q1 2008
Documentation					
	Proportion of TRaC and TRaC-M studies from 2005 to present w PSI/Washington Research MIS				
Asia			59%	72%	81%
Eastern Africa			14%	17%	24%
Latin America / Caribbean / E. Europe			18%	22%	22%
Southern Africa			8%	21%	58%
West and Central Africa			5%	7%	14%
Total			21%	27%	36%
Effectiveness					
	Proportion of countries with evidence of PSI impact with regard t				
Asia	21%	21%	21%	21%	21%
Eastern Africa	11%	11%	11%	11%	10%
Latin America / Caribbean / E. Europe	8%	8%	8%	8%	8%
Southern Africa	13%	13%	13%	13%	14%
West and Central Africa	40%	40%	40%	40%	46%
Total	21%	21%	21%	21%	21%
FoQus on Scales Capacity					
	Proportion of countries with a research manager who has attende				
Asia				0%	0%
Eastern Africa				78%	80%
Latin America / Caribbean / E. Europe				0%	0%
Southern Africa				88%	86%
West and Central Africa				73%	85%
Total				43%	45%
MAP: Phase One or Two Coverage					
	Proportion of PSI countries which have conducted a Phase One o				
Asia	50%	50%	58%	58%	58%
Eastern Africa	67%	67%	67%	67%	60%
Latin America / Caribbean / E. Europe	75%	75%	75%	75%	75%
Southern Africa	100%	100%	100%	100%	100%
West and Central Africa	53%	53%	53%	67%	54%
Total	66%	66%	68%	71%	67%

MAP: Phase Two Coverage	Proportion of PSI countries which have conducted a Phase Two M				
Asia	29%	29%	33%	42%	42%
Eastern Africa	22%	22%	33%	33%	30%
Latin America / Caribbean / E. Europe	75%	75%	75%	75%	75%
Southern Africa	63%	63%	63%	63%	71%
West and Central Africa	13%	13%	13%	13%	15%
Total	38%	38%	41%	43%	44%
SMRS Reports on Web					
	Total number of Social Marketing Research Series reports available on the Metrics page of the PSI website				
Asia	9	10	18	28	36
Eastern Africa	10	10	11	13	19
Latin America / Caribbean / E. Europe	8	10	11	11	12
Southern Africa	4	6	9	22	25
West and Central Africa	3	6	8	11	21
Total	34	42	57	85	113
Targeting					
	Proportion of PSI country programs which sell condoms that have resulted in changes in DALY parameters				
Asia		7%	7%	7%	7%
Eastern Africa		0%	0%	0%	30%
Latin America / Caribbean / E. Europe		0%	0%	0%	0%
Southern Africa		13%	13%	13%	43%
West and Central Africa		7%	7%	7%	8%
Total		5%	5%	5%	14%
TRaC: Two Round Country Coverage					
	Proportion of PSI countries with at least one risk group monitored in the PERForMance Review. Target: 80%				
Asia	86%	86%	86%	86%	86%
Eastern Africa	44%	44%	44%	44%	50%
Latin America / Caribbean / E. Europe	25%	25%	25%	25%	25%
Southern Africa	75%	75%	75%	75%	71%
West and Central Africa	60%	60%	60%	60%	69%
Total	59%	59%	59%	59%	61%
TRaC-M Coverage					
	Proportion of PSI countries which have conducted a TRaC-M study				
Asia	7%	7%	7%	7%	14%
Eastern Africa	0%	0%	0%	0%	10%
Latin America / Caribbean / E. Europe	8%	8%	8%	8%	8%
Southern Africa	0%	0%	0%	0%	0%

West and Central Africa	13%	13%	13%	13%	15%
Total	7%	7%	7%	7%	11%
FoQus on CD Coverage					
FoQus on CD Coverage	Proportion of PSI countries which have completed a FoQus Conc				
Asia / Eastern Europe / Pacific					
East Africa					
Latin America / Caribbean					
Southern Africa					
West and Central Africa					
Total					
MAP: Phase Three Coverage					
MAP: Phase Three Coverage	Proportion of PSI countries which have conducted a Phase Three				
Asia / Eastern Europe / Pacific					
East Africa					
Latin America / Caribbean					
Southern Africa					
West and Central Africa					
Total					
Targeting (last 3 years)					
Targeting (last 3 years)	Minimum Standard: All interventions spending \$200,000 or more must supply that information at least every three years. Interventions must have submitted collected targeting data and confirmed by the Me				
Asia / Eastern Europe / Pacific					
East Africa					
Latin America / Caribbean					
Southern Africa					
West and Central Africa					
Total					
BCC DALYs					
BCC DALYs	Minimum Standard: All interventions aiming at preventing HIV transmission must collect data with suggested questions in at least one round of data with suggested questions in at least one round. Definition: Proportion of PSI countries which conduct behavior change communication that have provided evidence of change in sexual partner, number of sexual contacts, condom use, needle/syringe use based on pre- and post-intervention TRaC data.				
Asia / Eastern Europe / Pacific					
East Africa					
Latin America / Caribbean					
Southern Africa					
West and Central Africa					
Total					

rd

Q2 2008	Q3 2008	Q4 2008	Q1 2009

with an SMRS report on the website and complete documentation in the

66%	65%	66%	56%
25%	32%	42%	55%
9%	52%	58%	53%
53%	53%	43%	52%
25%	47%	49%	51%
39%	51%	60%	52%

to any behavior. Source: PERForMance Review. **Target: 100%**

25%	25%	25%	50%
10%	10%	10%	40%
0%	0%	0%	30%
14%	14%	14%	57%
43%	43%	43%	36%
21%	21%	21%	42%

d a FoQus on Scales training

0%	0%	0%	
80%	80%	80%	
0%	0%	0%	
86%	86%	86%	
79%	79%	79%	
44%	44%	44%	

r Phase Two MAP study. **Target: 100%**

57%	57%	57%	57%
60%	60%	60%	60%
90%	90%	90%	90%
100%	100%	100%	100%
50%	64%	64%	64%
67%	71%	71%	71%

MAP study. Target: 100%

36%	36%	36%	36%
30%	30%	30%	30%
90%	90%	90%	90%
71%	71%	71%	86%
21%	36%	36%	43%
45%	49%	49%	53%

able for download on the Research &

50	57	59	
29	32	35	
10	20	25	
34	34	40	
30	49	52	
153	192	211	

e reported targeting information resulting

6%	6%	33%	
30%	30%	60%	
30%	30%	67%	
43%	43%	57%	
7%	7%	38%	
19%	19%	52%	

d with two rounds of data. Source:

88%	88%	88%	
50%	50%	50%	
10%	60%	60%	
86%	86%	86%	
64%	64%	64%	
61%	70%	70%	

udy.

13%	13%	13%	13%
10%	10%	10%	10%
10%	10%	10%	10%
0%	0%	0%	14%

14%	14%	14%	14%
11%	11%	11%	12%
Concept Development study.			
0%	0%	0%	19%
0%	0%	0%	30%
10%	20%	20%	20%
0%	0%	0%	29%
7%	7%	7%	14%
4%	5%	5%	21%

MAP study.

			0%
			0%
			0%
			0%
			7%
			2%

Frequency per year for which "targeting" information is useful for the DALY

Definition: Proportion of PSI countries that sell distribute condoms that are in DC.

			6%
			30%
			40%
			71%
			21%
			28%

Transmission (heterosexual, homosexual, or injection drug use) must supply at least every three years.

Communication campaign to prevent HIV transmission through heterosexual, desired change in behaviors of interest (i.e., being abstinent, number of syringe sharing, initiation of injection, etc.) as a result of PSI intervention

			2
			1
			0
			0
			0
			3